

## Overview and Outlook

Abhay Ashtekar

*Center for Gravitational Physics and Geometry*

*Pennsylvania State University, University Park, PA 16802-6300*

*When you do some mountaineering...you sometimes...want to climb to some peak but there is fog everywhere...you have your map or some other indication where you probably have to go and still you are completely lost in the fog. Then...all of a sudden you see quite vaguely in the fog, just a few minute things from which you say, "Oh, this is the rock I want." In the very moment that you have seen that, then the whole picture changes completely, because although you still don't know whether you will make the rock, nevertheless for a moment you say, "... Now I know where I am; I have to go closer to that and then I will certainly find the way to go ..."*

(Werner Heisenberg, 1963)

### 1. Introduction

The problem of quantum gravity is an old one and over the course of time several distinct lines of thought have evolved. However, for several decades, there was very little communication between the two main communities in this area: particle physicists and gravitation theorists. Indeed, there was a lack of agreement on even what the key problems are. By and large, particle physics approaches focused on perturbative techniques. The space-time metric was split into two parts:  $g_{\mu\nu} = \eta_{\mu\nu} + Gh_{\mu\nu}$ ,  $\eta_{\mu\nu}$  being regarded as a flat kinematic piece,  $h_{\mu\nu}$  being assigned the role of the dynamical variable and Newton's constant  $G$  playing the role of the coupling constant. The field  $h_{\mu\nu}$  was then quantized on the  $\eta_{\mu\nu}$ -background and perturbative techniques that had been so successful in quantum electrodynamics were applied to the Einstein-Hilbert action. The key problems then were those of handling the infinities. The gravity community, on the other hand, felt that a central lesson of general relativity is that the space-time metric plays a dual role: it is important that one and the same mathematical object determine geometry *and* encode the physical gravitational field. From this perspective, an ad-hoc split of the metric goes against the very spirit of the theory and must be avoided. If one does not carry out the split, however, a theory of quantum gravity would be simultaneously a theory of quantum geometry and the notion of quantum geometry raises a variety of conceptual difficulties. If there is no background space-time geometry –but only a probability amplitude for various possibilities– how does one do physics? What does causality mean? What is time? What does dynamics mean? Gravity theorists focused on such conceptual issues. To simplify mathematics, they often truncated the theory by imposing various symmetry conditions

---

. To appear in the proceedings of the workshop on the Canonical Formalism in Classical and Quantum General Relativity, held at Bad-Honef, Germany in September 1993; H. Friedrich (ed), Springer-Verlag.

and thus avoided the field theoretic difficulties. Technically, the emphasis was on geometry rather than functional analysis. It is not that each community was completely unaware of the work of the other (although, by and large, neither had fully absorbed what the other side was saying). Rather, each side had its list of central problems and believed that once these issues were resolved, the remaining ones could be handled without much difficulty. To high energy theorists, the conceptual problems of relativists were perhaps analogous to the issues in foundations of quantum mechanics which they considered to be “unimportant for real physical predictions.” To relativists, the field theoretic difficulties of high energy physicists were technicalities which could be sorted out after the conceptual issues had been resolved. Even on the rare occasions when they got together, the two sides seemed to “talk past each other.”

Over the last decade, however, there has been a certain rapprochement of ideas on quantum gravity. Each side has become increasingly aware of the difficulties that were emphasized by the other. There are some agreements and increasing clarity. In the poetic imagery of Heisenberg, we have seen faint outlines of some of the rocks we want to climb.

Perhaps the most important among these developments are the following:

1. Recognition that non-perturbative methods are essential. There is a growing sentiment that, at a fundamental level, the theory should not involve fluctuations around a classical geometry. Indeed, there should be no background geometry or any background fields for that matter. The theory should be diffeomorphism invariant.
2. Acceptance that conceptual issues such as the problem of time will have to be handled satisfactorily. The unease with the standard measurement theory of quantum mechanics is brought to the forefront by the absence of a background, classical space-time. The interface of classical and quantum domains and the decoherence processes that make the world seem classical are receiving greater attention.
3. Agreement that the field theoretic divergences have to be faced squarely. There is growing awareness that although the mini-superspaces that feature in quantum cosmology are obviously interesting, from the perspective of full quantum gravity, they are essentially toy models. One has to learn to deal with the infinite number of degrees of freedom “honestly.” Time has come to give proper mathematical meaning to the formal equations such as the Wheeler-DeWitt equation. New ideas and mathematical techniques are needed since the issue of regulating quantum operators in absence of a background geometry has rarely been faced.
4. Reconciliation with the possibility that the “quantum geometry” governing the Planck scale may not at all resemble the Riemannian picture. It is likely that differential geometry, in the standard sense of the term, will itself be inadequate to capture physics. Some discrete structures are likely to emerge and new mathematical tools will be needed to handle them.

Thus, the goals of the two communities have moved closer.

These recognitions do not imply, however, that there is a general consensus on *how* all these problems are to be resolved. Thus, there are again many approaches. But I strongly support belief that this diversity in the lines of attack is very healthy. In a problem like quantum gravity, where directly relevant experimental data is scarce, it would be a grave error if everyone followed the same path. Indeed, the most promising way of enhancing

our chances at success is to increase the amount of variety. What is striking is that, in spite of the diversity of methods, some of the results are often qualitatively similar. For example, in the last few years, the fourth point I mentioned above has come up again and again in quite different contexts. The idea that the continuum picture itself is likely to break down is not new. For the first time, however, it is arising as a *concrete result* of calculations within well-defined, coherent schemes which were not explicitly constructed to obtain such a result. It arose from certain computer simulations of 4-dimensional Euclidean gravity, (see e.g. Agishtein & Migdal (1992) and Brüggmann & Marinari (1993)), from string theory (see e.g. Gross & Mende (1988), Amati et al (1990), Aspinwall (1993)), and, from canonical quantization of 4-dimensional general relativity (Ashtekar et al (1992)). The detailed pictures of the micro-structure of space-time that arise in these approaches are quite different at least at first sight. Nonetheless, there *are* certain similarities in the results; most of them are obtained by using genuinely non-perturbative techniques. The overall situation makes me believe that we have, so to say, entered a new era in the field of quantum gravity. Through the dense fog, we have caught a fleeting glimpse of the rocks that, we think, will bring us to the distant peak and, in the spirit of Heisenberg, various groups are charting their paths even though they still do not know if they will make it.

In this article, I will attempt to provide a broad overview of the canonical approach based on connections and loops and suggest some directions for future work. I should emphasize that this is not a systematic survey. I will not even attempt to cover all areas in which significant developments have occurred. Rather, I will concentrate on issues which are related to the topics covered by other speakers but which were not discussed in their talks in any detail. I should also emphasize that most of what I will say are personal perspectives which are not necessarily shared by others in the field. (For comprehensive reviews of the canonical approach based on connections and loops see other articles in this volume and, e.g., Ashtekar (1991,1992) and Pullin (1993).)

I begin in section 2 with an assessment of the canonical approach and, using 2+1-dimensional general relativity, illustrate the vision that underlies the program. In section 3, I will summarize some recent mathematical developments which were mentioned by Bernd Brüggmann (1994) and Renate Loll (1994) in their talks. These results have made it possible to define the loop transform rigorously for non-Abelian connections with an infinite number of degrees of freedom and they provide a basis for regulating and solving the quantum constraints. Section 4 illustrates how low energy (i.e. laboratory scale) physics can arise from non-perturbative quantum gravity and section 5 presents some promising directions for future work.

## 2. The canonical approach

This workshop as a whole was devoted to canonical methods in general relativity, with a particular emphasis on quantization. Therefore, in the first part of this section, I will present an assessment of the strengths and weaknesses of these methods in general terms, without restricting myself to the choice of a specific type of dynamical variables or to the details of how the canonical program is carried out. In the second part, I will present my own view of the general framework one can hope to arrive at in quantum gravity through the use of canonical methods.

## 2.1 An assessment

At the mathematical as well as conceptual level, the quantum theory we best understand is the non-relativistic quantum mechanics of point particles. In this case, the canonical approach provides the “royal road” to quantization. One can even formulate the fundamental kinematic problem as that of obtaining the appropriate representation of the basic canonical commutation relations. The problem of dynamics is then that of defining the Hamiltonian operator and of fully understanding its action on states. Note that the path integral method is *not* a substitute for canonical quantization. It is rather an alternative –and, in the case of scattering problems, a powerful– method of tackling the issue of quantum dynamics. What it provides is the transition amplitudes  $\langle \vec{x}', t' | \vec{x}, t \rangle$  for the particle to go from the position  $\vec{x}$  at time  $t$  to the position  $\vec{x}'$  at time  $t'$ . The full quantum mechanical results, however, involve calculating the amplitude for a transition from one *physically realizable* quantum state to another –say, from a state  $\Psi_0(\vec{x})$  with angular momentum 0 to a state  $\Psi_m(\vec{x})$  with angular momentum  $m$ . And, to obtain these, we need to know the Hilbert space of states, the action of physically interesting operators and the eigenfunctions and eigenvalues of these operators. These have to be supplied externally –typically through canonical quantization. It is for this reason that most courses on non-relativistic quantum mechanics focus on the canonical method.

Coming to field theory, the canonical approach is again the oldest coherent method to tackle the problem of quantization. It is deeply rooted in the Heisenberg uncertainty relation and hence in the fundamental *physical* principles of quantum mechanics. Why is it then that no modern course in field theory develops the subject within the canonical method? The main reasons, I believe, are the following. First, the approach lacks *manifest* covariance. Second, it is not well-suited to the use of Feynman diagrams and other perturbative techniques. The first drawback arises from obvious reasons: The basic canonically conjugate fields satisfy equal time commutation relations and are therefore defined on a Cauchy slice, whence the 4-dimensional covariance is broken at the outset. The origin of the second drawback lies in the fact that the quantum states in the canonical approach arise as suitable functionals of fields and this representation is quite inconvenient if one wants to work with particle states. In perturbative field theory, on the other hand, the emphasis is on particles –real and virtual– and the powerful machinery of Feynman diagrams is tuned to the particle picture. Furthermore, each diagram represents a *space-time* process. The exchange of virtual particles, particles moving forward in time and the anti-particles moving backwards, real particles scattering off each other –each of these processes involves the passage of time and is a representation of a 4-dimensional, space-time integral. It is in principle possible but in practice cumbersome to capture all this in the canonical framework. It is hard to prove renormalizability. It is *much* easier to forego the descriptions in terms of functionals of 3-dimensional fields and consider instead the Fock spaces of particle states. Indeed, this is hardly surprising. It is always the case that calculations simplify when one tailors the framework to the dynamics in question. In the perturbative treatment of scattering theory, then, simplifications should occur when we use eigenstates of the asymptotic, free Hamiltonian. And these are precisely the particle states in the Fock space.

One might adopt the viewpoint that from the standpoint of what is true “in principle,”

both the drawbacks stem from aesthetic considerations. Indeed, although the procedure is not manifestly covariant, it is nonetheless true that, in the final picture, the quantum field theory one obtains, say in the case of a free Maxwell field, is completely equivalent to the more familiar one featuring the Fock space of photons. In the canonical approach, the states are represented by (gauge invariant) functionals  $\Psi(A)$  of the vector potential  $A_a(\vec{x})$  on a 3-dimensional plane (or, more generally, a Cauchy surface) in Minkowski space. This description breaks the manifest covariance. However, the Poincaré group *is* unitarily implemented. The vacuum state *is* Poincaré invariant. The spectrum of the (free) 4-momentum operator *is* causal and future-directed, and so on. Similarly, as we noted already, using the equivalence with the Fock representation, one can translate, step by step, any perturbative calculation in, say, quantum electrodynamics to the canonical language. This is all true. But the price one would pay would be substantial. It would be analogous to –and *much* more complicated than– insisting on using Cartesian coordinates in a problem where the Hamiltonian has manifest spherical symmetry. Not only would it make calculations enormously more difficult but one would also lose all sorts of *physical* insights. This is not to say that canonical quantization has no place in field theory. It does. Most text books begin the discussion with canonical quantization perhaps because the underlying basic structure is clearest in this framework. Similarly, at the odd place when one is not sure of the measure in the path integral, one returns to the canonical picture and “derives the correct measure” starting with the Liouville form on the phase space. However, for the actual calculations of scattering cross sections or probing general properties such as renormalizability, the language of particles and the 4-dimensional pictures are obviously better-suited.

What about general relativity? The canonical approach has the great advantage that it does not require us to introduce a background metric. It is well suited for a non-perturbative treatment. The final Hamiltonian framework does have a number of features not encountered in Minkowskian field theories. However, in view of the profound conceptual differences between these theories and general relativity, the emergence of such features is but to be expected. In particular, much of the dynamical information of the theory is now contained in constraints. Indeed, as discussed by Robert Beig (1994), in the spatially compact case, the Hamiltonian vanishes identically on the constraint surface, i.e., on the physical states of the classical theory. In quantum theory then, to begin with, there is no Hamiltonian, no time and no evolution. One only has physical states –the solutions to quantum constraints. Yet, it may be possible to “extract” dynamics from these solutions by identifying a suitable physical variable as an internal clock. Indeed, this has already been done in 2+1-dimensional gravity (see e.g. Carlip (1993)). Thus, at least in principle, the approach provides us with a well-defined, precise strategy that is sufficiently sophisticated to extract physical information at the quantum level in spite of the absence of a background geometry. This is its greatest strength.

How does the approach fare with respect to the two key difficulties it faces in Minkowskian quantum field theories? In quantum gravity, the scattering cross sections are not our *prime* concerns. Since there is no background space-time metric, notions such as particle states –and gravitons in particular– which are tied to the Poincaré group would presumably be only approximate concepts. The fact that quantum general relativity (and

various modifications thereof) fails to provide us with a consistent, local quantum field theory perturbatively also indicates that the fundamental theory should not be formulated in terms of these particle states. The whole imagery of processes mediated by particles moving forward and backward in time is probably inappropriate. So, the second reason for abandoning the canonical method now loses its force. Indeed, the key question now is if the quantum dynamics of the gravitational field can be made simple in an appropriate representation within the canonical scheme. And, as we have seen in this workshop, there are indeed strong indications that the loop representation is well-suited for this purpose (see e.g. Brügmann (1994)).

The lack of manifest covariance of the canonical scheme, however, is still with us. And indeed it is now a *much more* serious issue. In Minkowskian field theories the issue was an aesthetic one. While the canonical procedure violates *manifest* covariance, it *is* covariant; the Poincaré group *is* unitarily implemented. We can, if we wish, describe dynamics in the space-time picture. In quantum gravity this is no longer the case. In classical general relativity, it is the space-time geometry that is the dynamical variable. A space-time represents a possible history –analogous to a trajectory in particle mechanics. Just as in non-relativistic quantum mechanics the particle trajectories have no basic role to play in the final quantum description, one would expect that in full canonical quantum gravity, space-times would have no distinguished place. There will presumably exist some special (“semi-classical”) states which can be approximated by 4-dimensional space-times. Given any such state, one would be able to speak of approximate 4-dimensional covariance in an effective theory which ignores large quantum fluctuations away from that state. At a fundamental level, however, there would not be a 4-dimensional geometric entity to replace the classical space-time. And if there is no such entity, how can one even speak of 4-dimensional covariance? It does seem that there is a branching of ways here. Furthermore, it is the qualitative features of the canonical approach that force this branching; it is not tied to the use of specific variables.

To preserve the space-time covariance, it is tempting to choose the path integral approach. Even though in the finished picture there is no preferred space-time, one does have an underlying 4-manifold and the basic objects that one calculates with are 4-metrics. And the diffeomorphism group does act on the space of these metrics. Since the Einstein-Hilbert action is diffeomorphism invariant, one would expect to get diffeomorphism invariant answers for (the correctly formulated) physical questions. Thus, the strategy seems attractive. There are, however, three problems.

First, since path integrals in full quantum gravity are only formal expressions, everything one does *from the very beginning* involves formal manipulations; the level of mathematical precision leaves much to be desired. In practice, one generally proceeds by making some “approximations” which mimic the successful strategies in other field theories. However, it is not at all obvious that these strategies can be taken over to gravity in a meaningful fashion. Indeed, in Minkowskian field theories such as quantum electrodynamics, the only way we know how to make sense of path integrals is through perturbation theory. In fact, the power of the method lies in the fact that it captures the enormous information in the perturbation expansions very succinctly. In a theory that does not exist perturbatively, we are at a loss. It would be surprising indeed if all of the approximation

methods that are useful in the perturbative context will continue to be useful in quantum gravity. A much more sophisticated approach is needed. One might imagine using the techniques developed in rigorous quantum field theory. However, in constructive quantum field theories, path integrals are defined in the Euclidean regime and the physical, Lorentzian Green's functions are obtained through a wick rotation. We all know that, in gravity, this simple route is not available.

The second problem is that while the method is manifestly covariant in the sense of classical general relativity, it lacks *quantum covariance* even in the case of particle dynamics. Let me explain this point in some detail since many of the readers may be unfamiliar with the issue. In the sum over histories approach, one fixes, at the outset, a preferred configuration space—all histories are to be trajectories in this space. In quantum mechanics, this corresponds to fixing a representation (such as the position or the momentum) of the observable algebra. Now, one of the technically powerful and, I feel, conceptually deep features of quantum physics is Dirac's transformation theory<sup>†</sup> which establishes the covariance of the *quantum* theory under the change of representation. This covariance is manifest in the canonical approach. In the path integral approach, not only it is not manifest but it is often cumbersome to incorporate. Consequently, just as it is often hard to phrase "space-time questions" in the canonical approach, it is hard to phrase questions which involve different representations in the path integral approach (unless of course one passes through the canonical approach and makes use of the Dirac transformation theory). Consider for example an harmonic oscillator. If one wants to ask for the probability for the particle which passes through the interval  $\Delta_0$  of the position space at time  $t_0$  and the interval  $\Delta_1$  at time  $t_1$  to end up in an interval  $\Delta_2$  at time  $t_2$ , we immediately know we should integrate over the set of paths which pass through the given intervals at the given times. Now suppose we change the question somewhat by replacing the condition at the intermediate time  $t_1$  and ask instead that, at that time, the particle have an energy  $\hbar\omega(n + \frac{1}{2})$  for a fixed  $n$ . Now, it is no longer obvious what paths to consider. More generally, paths or histories are trajectories in *some* configuration space (i.e. a Lagrangian sub-manifold of the phase space) and path integrals can easily cope with questions that refer to that fixed sub-space. In quantum theory, by contrast, the domain space of wave functions may have nothing to do with the classical phase space. For example, in the case of the harmonic oscillator, we could use the energy representation in which the quantum states are functions  $\Psi(n) = \langle n | \Psi \rangle$  of an integer which (do not constitute a Lagrangian sub-manifold of the phase space and therefore) can not serve as a classical configuration space. Path integrals are not well-adapted to such representations. But it may well be that it is precisely such representations that are best suited to incorporate quantum dynamics. The loop representation, in particular, falls in this category.

The last problem is the one I already mentioned in the beginning of this sub-section. Path integrals provide transition amplitudes but not the kinematical structure; it has to be supplied from outside. In a real sense, therefore, the path integral and the canonical approaches generally complement each other. In quantum gravity, it is possible that

---

<sup>†</sup> about which Dirac said in 1977: "I think that is the piece of work which has most pleased me of all the works that I have done in my life...The transformation theory became my darling." (Pais (1987))

ultimately the two will be used together so that the framework has both classical and quantum covariance. The dynamics coded in the quantum constraints in the canonical approach could perhaps be re-interpreted in terms of mathematically well-defined path integrals to demonstrate that there is a well-defined sense in which the theory enjoys space-time covariance. Such an interpretation would be especially helpful in the analysis of whether topology changes occur in quantum gravity and, if so, whether they have physically significant ramifications in the low energy regime. The full quantum covariance could be established by switching to the canonical framework.

It may also turn out, however, that in the gravitational case, canonical quantization can not be reconciled with path integrals in the full quantum theory. As the discussion in the next sub-section indicates, even the “spatial” 3-manifold could be a secondary construct. At the fundamental level, there may only be discrete structures and combinatorial operations. In such a scenario, the continuum picture and objects such as manifolds may arise as useful mathematical constructs only in semi-classical physics. And the theory would have the desired covariance only in these regimes. It may be that the predictions of the two methods agree in such regimes but in the fully Planckian domain, the two theories are quite different.

To summarize then, the canonical methods are well adapted to non-perturbative treatments of quantum gravity. However, there appears to be a fundamental tension between the quantum gravity scenarios that are natural to canonical approaches and space-time pictures that we are so accustomed to in the classical regime. Whether this tension is real or only apparent is not yet clear. Technical progress, particularly in the path integral approach, would be of great help to settle this issue.

## 2.2 An underlying vision

In this workshop, Thomas Thiemann discussed 2+1-dimensional gravity in some detail and Hermann Nicolai presented an extension (due to de Wit, Matschull and himself (1993)) of those results to give an elegant clarification of the issue of the “size” of the space of physical states in supergravity. I would now like to use 2+1 gravity for a different purpose: to illustrate my own expectations of non-perturbative quantum gravity in 3+1 dimensions. (For details on 2+1-dimensional gravity, in addition to these proceedings, see, e.g., Carlip (1990, 1993), and Ashtekar (1991), chapter 17.)

Let me begin with a brief historical detour. Conceptually –and, in certain respects, also technically– 2+1-dimensional general relativity is very similar to the 3+1-dimensional theory. There is no background structure; the theory is diffeomorphism invariant. One and the same object –the space-time metric  $g_{\mu\nu}$ – determines the geometry *and* encodes the gravitational field. In the canonical description, the Hamiltonian again vanishes in the spatially compact case and the dynamics is driven by first class constraints. Furthermore, in geometrodynamics, the general structure of the constraint algebra is the same as in the 3+1 theory (see, e.g., Beig 1994); it is not a true Lie algebra. Consequently, for a number of years, it was believed that a non-perturbative, canonical quantization of this theory is as difficult as that of the 3+1-dimensional theory. In particular, not a single solution to the Wheeler-DeWitt equation was known. In the perturbative treatment, one finds that Newton’s constant is again dimensionful (with dimension  $[L]$ , thus a *positive* power of length in the  $\hbar = c = 1$  units) and simple power counting arguments suggested that the



theory is not renormalizable. Therefore, there appeared several papers in the literature saying that 2+1-dimensional general relativity is as intractable as the 3+1-dimensional theory.

This would have been somewhat puzzling since in 2+1-dimensions there are no local degrees of freedom; all solutions to the Einstein field equations are flat. Indeed, the conclusions of these papers turned out to be wrong. There are no divergences. Using connections as basic variables rather than metrics, the quantum theory can in fact be solved exactly in the spatially compact case for any genus. (Achúcarro & Townsend (1986), Witten (1988). See also Ashtekar et al (1989) and Nelson & Regge (1989)). Since then, the theory has also been solved in the metric representation in the case when the spatial topology is that of a 2-torus (Moncrief 1989, Hosoya & Nakao (1990)) and work is in progress on the higher genus case. One of the lessons that one can draw from this work is that connections and holonomies are better suited to the mathematics of the quantum theory than metrics and light cones. It is *not* that the theory can not be solved in the metric picture; as I just remarked, this has already been achieved in the 2-torus topology. Rather, connections seem to be better adapted to deal with the quantum constraints and to write down Dirac observables. Variables which are most useful to the macro-physics are not the ones which make the micro-physics most transparent.

There are several equivalent ways of quantizing 2+1 gravity on a torus (see, e.g., Carlip 1993). The variety of representations that result provide us with a rich example of the Dirac transformation theory in action. Here, I will consider three and argue that each clarifies and emphasizes a different aspect of quantum gravity. At first sight, it appears that each representation provides its own, distinct picture of reality. And yet, thanks to the Dirac transformation theory, they are all equivalent.

First, there is the loop representation. Here, quantum states are functions of homotopy classes of loops on the spatial 2-manifold; the domain space of quantum states is thus *discrete*. In the case of a torus, each homotopy class is represented by a pair of integers  $(n_1, n_2)$  which tell us how many times the loop winds around the two generators of the homotopy group. Hence, in this case, the quantum states  $\Psi(n_1, n_2)$  are functions just of two integers. The basic observables of the theory,  $T^0(n_1, n_2)$  and  $T^1(n_1, n_2)$ , are also labelled by two integers and their action on states just shifts the arguments of the wave functions. Thus, the whole mathematical structure is *combinatorial*. There is no space, no time, no continuum. Hence there is no diffeomorphism group to implement and no issue of space-time covariance to worry about. I would like to regard this as the “fundamental description.” Now, we learnt from Thomas Thiemann’s talk (1994) that this description is completely equivalent to that in terms of connections on a 2-torus<sup>†</sup>. Put differently, suppose we knew nothing about general relativity but were provided with just the combinatorial description given above. Staring at this description, a clever young student could have realized that the description would become “nice and geometric” if one were to introduce a

---

<sup>†</sup> The two are related by the loop transform. A certain amount of caution is needed in the definition of the loop transform however because the space on which one integrates—the moduli space of flat connections—has non-compact connected components on which the Wilson loop functions are unbounded (Marolf 1993, Ashtekar & Loll (1993)). This important technical point was overlooked in the earlier treatment (Ashtekar et al 1989).

“fictitious, mathematical” 2-manifold, namely a 2-torus, and regard the arguments  $(n_1, n_2)$  of quantum states as labels for the homotopy classes of closed loop on this torus. Excitedly, she tells us that, once this is done, one can equally consider the quantum states of two real numbers  $(a_1, a_2)$  which can be regarded as labels for gauge equivalent flat connections on the torus.. After working in this picture for a while, we realize that the theory can be regarded as quantization of an interesting classical theory which features connections on a 3-dimensional manifold, obtained by taking the product of our fictitious torus with a fictitious real line. To our astonishment, we then notice that the theory has the peculiar property of being diffeomorphism invariant. Furthermore, we notice that it can be thought of as a theory of metrics of signature  $-++$  and the connection which we discovered earlier is the spin connection of this metric. We can then go on and study the light cones and causal properties of this classical theory and be surprised that all this rich and unexpected structure arises in the classical limit. We discover that, in this classical approximation, the 3-manifold can be thought of as “space-time” in which things live. Intrigued by this notion of time and dynamics, we return to the quantum theory and realize that with a little bit of work, we can construct a third representation in which quantum states are functions of *three* variables subject to a differential equation which tells us how the state changes or “evolves” as we increase the third variable. We realize that variable could be interpreted as an “internal clock.”

Thus, we have three different quantum pictures. The “fundamental” description is combinatorial. The second, in terms of connections, is a “timeless” description. We can interpret the 2-manifold as “space” and see that the spatial diffeomorphisms are unitarily implemented. But there is no time; we have a “frozen formalism” a la Bergmann and Komar. Finally, in the third description, we reintroduce time through an internal clock. Now the description resembles that of a parametrized particle. We are on familiar ground.

My hope is that we will arrive at a similar description in the 3+1 theory. There will be several equivalent representations, each illuminating its favorite facets of quantum gravity. Thanks to the Dirac transformation theory, the numerous expectations we have of the theory –which often seem even to contradict each other– will in fact be compatible. There is already some evidence in favor of this scenario. The loop and connection representations are qualitatively similar to those in the 2+1-dimensional case. (In fact, the emphasis on connections in the 2+1-dimensional theory was motivated, in the first place, by the connection dynamics formulation of the 3+1 theory.) In the loop representation, the diffeomorphism constraint leads us to knot classes and hence to combinatorial operations. The connection representation will lead us directly to a timeless, frozen formalism. And as we will see in section 4, in the weak field truncation, we can identify one of the components of the connection as an internal clock and recover quantum dynamics from the Hamiltonian constraint. Thus, we do have a number of pointers. Of course, whether this scenario is borne out in detail remains to be seen. In my view, the weakest point may be the last one; we may be able to isolate internal clocks only under suitable approximations.

### 3. Recent mathematical developments

It is obvious from the articles by Brügmann, Loll and Thiemann that the loop representation plays an important role in this approach to quantum gravity. It is therefore

important to put this representation and results obtained therefrom on a sound mathematical footing. It is worth emphasizing that this is not just an abstract desire for rigor. In a real sense, in non-perturbative quantum gravity, we are entering a completely uncharted territory and the mathematical techniques required are quite new. In most of the work to date, one has proceeded by making convenient choices –e.g., of the regularity conditions on loop states, of factor-ordering and of limiting procedures used to regulate operators– as the need arose. These choices do seem “reasonable” and there *is* an overall coherence to the entire scheme. However, it is not clear that there aren’t other choices that are equally plausible and that the results will not change if these choices were made. There are no uniqueness results. And none will come by unless one elevates the degree of mathematical precision. In a sense, this is analogous to the situation in interacting quantum field theory in Minkowski space, where there is again no a priori justification for using the Fock representation and the associated regularity conditions on states and operators. (In fact, in view of Haag’s theorem (see, e.g., Streater & Wightman (1964)), there are good mathematical reasons for *not* using it!). However, in that case, over the years one has acquired a great deal of theoretical experience and, more importantly, there is enormous experimental data which supports the strategy as a working hypothesis. In non-perturbative gravity, on the other hand, one lacks both and the issue of control over the assumptions and uniqueness results become much more important.

Fortunately, over the last two years several mathematical developments have occurred which have the potential of improving the situation dramatically. The finished and published results deal only with real  $SU(n)$  or  $U(n)$  connections. However, work is in progress to extend these results to the complex-valued connections that one encounters in gravity and it appears that the key results which enable one to define the loop transform and the loop representation will go over to that case as well<sup>†</sup>. For brevity and precision, in this section, I will restrict myself to  $SU(2)$  connections and report on the results that have been already obtained.

To define the loop transform, we need an integration theory on the space  $\mathcal{A}/\mathcal{G}$  of connections modulo gauge transformations. In the 2+1-dimensional case, this problem is easy to solve because the (appropriate components of the) moduli space of *flat*  $SU(1,1)$  connections can be naturally given the structure of a finite dimensional symplectic manifold (see e.g. Ashtekar (1991), chapter 17); one can simply use the Liouville volume element to evaluate integrals. Thus, although the domain space of quantum states is non-linear,

---

<sup>†</sup> The idea is to let the gauge group be  $SU(2)$  –as is needed for the triads  $E_i^a$  to be real– but to allow complex-valued connections  $A_a^i$  (which take values in the Lie-algebra of  $\mathbb{C}SU(2)$ , complexified  $SU(2)$ ). Thus, we are using the original Hamiltonian framework in which  $A_a^i$  is regarded as a complex-valued coordinate on the real phase space of general relativity. One then uses appropriate extensions of the techniques outlined below to develop calculus on the resulting  $\mathcal{A}/\mathcal{G}$ . In particular, one can give a precise meaning to the notion of holomorphic functionals used in the connection representation. We expect that the appropriate parts of the integration theory will also go through and enable us to define the loop transform rigorously, although here the work is still incomplete. If all our expectations are borne out, we would also have incorporated the reality conditions at the kinematical level, i.e. prior to the imposition of constraints.

the integration theory is simple because the space is *finite* dimensional. In 3+1 space-time dimensions, the situation is again simple for the case of *linearized gravity* –the theory of free gravitons in Minkowski space. This theory can be cast in the language of connections (see, e.g., Ashtekar (1991), chapter 11). Integration theory is again well-developed; the domain space is now linear and one can simply use the Gaussian measure as in free field theories in Minkowski space. Thus, in this case, in spite of the presence of an infinite number of degrees of freedom, the integration theory is straightforward because of the underlying *linearity*. The problem is significantly more difficult if  $\mathcal{A}/\mathcal{G}$  is *both* non-linear *and* infinite dimensional.

Let us now discuss this case. Let us begin by fixing an analytic 3-manifold  $\Sigma$  which will of course represent a Cauchy surface in space-times to be considered. We will consider  $SU(2)$  connections on  $\Sigma$ . Since any  $SU(2)$  bundle over a 3-manifold is trivial, we can represent any connection by a Lie-algebra valued 1-form  $A_a^i$  on  $\Sigma$ , where  $a$  is the spatial index and  $i$ , the internal<sup>†</sup>. Denote by  $\mathcal{A}$  the space of smooth (say  $C^2$ )  $SU(2)$ -connections equipped with one of the standard (Sobolev) topologies (see, e.g. Mitter and Viallet (1981)).  $\mathcal{A}$  has the structure of an affine space. However, what is of direct interest to us is the space  $\mathcal{A}/\mathcal{G}$  obtained by taking the quotient of  $\mathcal{A}$  by  $(C^3)$  local gauge transformations. In this quotient construction, the affine structure is lost;  $\mathcal{A}/\mathcal{G}$  is a genuinely non-linear space with complicated topology.

To define the integration theory, we will adopt an algebraic approach. To see the main idea, consider integrals  $\int f d\mu$  of continuous functions  $f$  on a compact, Hausdorff space  $S$  with respect to a regular (but not necessarily strictly positive) measure  $d\mu$ . The space of continuous functions has the structure of an Abelian  $C^*$ -algebra with identity and the map  $f \rightarrow \langle f \rangle := \int f d\mu$  is a functional on the algebra satisfying the following two properties: i) positivity (which should really be called non-negativity), i.e.  $\langle \bar{f}f \rangle \geq 0$  for any element  $f$  in the algebra; and, ii) linearity, i.e.  $\langle f + \lambda g \rangle = \langle f \rangle + \lambda \langle g \rangle$  for all elements  $f$  and  $g$  of the algebra and complex numbers  $\lambda$ . Thus, given a measure, we acquire a positive linear functional on the given Abelian  $C^*$ -algebra. Now, standard theorems ensure us that *every* regular measure on  $S$  arises in this manner. The idea now is to use this 1-1 correspondence to define measures on the space of connections modulo gauge transformations.

Let us begin by constructing an appropriate sub-algebra of the Abelian  $C^*$ -algebra of continuous, bounded functions on  $\mathcal{A}/\mathcal{G}$ . This is to be the algebra of configuration variables. Therefore, the obvious strategy is to use the Wilson loop functionals. Following the procedure and notation used by other speakers, given any closed loop  $\alpha$  on the 3-manifold  $\Sigma$ , let us define the Wilson-loop functional  $T_\alpha$  on  $\mathcal{A}/\mathcal{G}$ :

$$T_\alpha(A) := \frac{1}{2} \text{Tr } \mathcal{P} \exp G \oint_\alpha A \cdot dl, \quad (3.1)$$

---

<sup>†</sup> Results reported in this section for which explicit references are not provided are all taken from Ashtekar and Isham (1992) and Ashtekar and Lewandowski (1993a,b). The last two papers and those by Baez (1993a,b) contain significant generalizations which include allowing more general gauge groups, allowing the manifold  $\Sigma$  to be of arbitrary dimension and allowing the connections to live in non-trivial bundles.

where the trace is taken in the fundamental representation of  $SU(2)$  and the Newton's constant  $G$  appears because, in general relativity, it is  $GA_a^i$  that has the dimensions of a connection; in gauge theories, of course, this factor would be absent. For technical reasons, we will have to restrict ourselves to piecewise analytic loops  $\alpha$ . (This is why we needed  $\Sigma$  to be analytic. Note that the loops need not be smooth; they can have kinks and self-intersections but only at a finite number of points.) As explained in detail by Brüggmann and Loll, due to  $SU(2)$  trace identities, the product of any two Wilson-loop functionals can be expressed as a sum of other Wilson loop functionals. Therefore, the vector space generated by finite complex-linear combinations of these functions has the structure of a  $\star$ -algebra (where the  $\star$ -operation is just complex-conjugation). The functionals  $T_\alpha$  are all bounded (between  $-1$  and  $1$ ). Hence, the sup-norm (over  $\mathcal{A}/\mathcal{G}$ ) is well-defined and we can take the completion to obtain a  $C^*$ -algebra. We will call it the *holonomy  $C^*$ -algebra* and denote it by  $\mathcal{HA}$ . Elements of  $\mathcal{HA}$  are to be thought of as the configuration variables of the theory.

Since  $\mathcal{HA}$  is an Abelian  $C^*$ -algebra with identity, we can apply a standard theorem due to Gel'fand and Naimark to conclude that  $\mathcal{HA}$  is isomorphic with the  $C^*$ -algebra of all continuous functions on a compact, Hausdorff space  $sp(\mathcal{HA})$ , the spectrum of the given  $C^*$ -algebra  $\mathcal{HA}$ . The spectrum itself can be recovered from the algebra  $\mathcal{HA}$  directly; its points are homomorphisms from  $\mathcal{HA}$  to the  $\star$ -algebra of complex numbers. Now, we know that the elements of  $\mathcal{HA}$  suffice to separate points of  $\mathcal{A}/\mathcal{G}$ ; given any two elements of  $\mathcal{A}/\mathcal{G}$  there is at least one loop  $\alpha$  such that  $T_\alpha$  takes different values on the two elements. Therefore it follows from the Gel'fand-Naimark theorem that  $\mathcal{A}/\mathcal{G}$  is *densely embedded* in  $sp(\mathcal{HA})$ . To emphasize this point, from now on, we will denote the spectrum by  $\overline{\mathcal{A}/\mathcal{G}}$  and regard it as a completion of  $\mathcal{A}/\mathcal{G}$  (in the Gel'fand topology). Integration theory will be defined on  $\overline{\mathcal{A}/\mathcal{G}}$  rather than  $\mathcal{A}/\mathcal{G}$ . This enlargement is in accordance with the common occurrence in quantum field theory: while the classical configuration (or phase) space may contain only smooth fields (typically taken to belong to be the Schwartz space), the domain space of quantum states is a completion of this space in an appropriate topology (the space of distributions). From the remarks I made above, it is clear that regular measures on  $\overline{\mathcal{A}/\mathcal{G}}$  will correspond to positive linear functions on the holonomy algebra  $\mathcal{HA}$ . This is the general strategy we will follow.

However, a key difficulty with the use of the Gel'fand theory is that one generally has relatively little control on the structure of the spectrum. A reasonable degree of control is essential since (at the kinematical level) the quantum states in the connection representation are to be functions  $\Psi(\bar{A})$  on  $\overline{\mathcal{A}/\mathcal{G}}$ . In the present case, we are fortunate: a simple and complete characterization of the spectrum is available. To present it, I first need to introduce a key definition. Fix a base point  $x_o$  in the 3-manifold  $\Sigma$  and regard two (piecewise analytic) closed loops  $\alpha$  and  $\alpha'$  to be equivalent if the holonomy of any connection in  $\mathcal{A}$ , evaluated at  $x_o$ , around  $\alpha$  is the same as that around  $\alpha'$ . We will call each equivalence class (a holonomically equivalent loop or) a *hoop* and denote the hoop to which a loop  $\alpha$  belongs by  $\tilde{\alpha}$ . For example,  $\alpha$  and  $\alpha'$  define the same hoop if they differ by a reparametrization or by a line segment which is immediately re-traced. (For piecewise analytic loops and  $SU(n)$  connections, one can show that these two are the most general operations; two loops define the same hoop if and only if they are related by a combination

of reparametrizations and retracings.) The set of hoops has, naturally, the structure of a group. We will call it the *hoop group* and denote it by  $\mathcal{HG}$ . In terms of this group, we can now present a simple characterization of the Gel'fand spectrum  $\overline{\mathcal{A}/\mathcal{G}}$ :

*Every homomorphism  $\hat{H}$  from the hoop group  $\mathcal{HG}$  to the gauge group  $SU(2)$  defines an element  $\bar{A}$  of the spectrum  $\overline{\mathcal{A}/\mathcal{G}}$  and every  $\bar{A}$  in the spectrum defines a homomorphism  $\hat{H}$  such that  $\bar{A}(\tilde{\alpha}) = \frac{1}{2} \text{Tr } \hat{H}(\tilde{\alpha})$ . This is a 1-1 correspondence modulo the trivial ambiguity that homomorphisms  $\hat{H}$  and  $g^{-1} \cdot \hat{H} \cdot g$  define the same element  $\bar{A}$  of the spectrum.*

Clearly, every regular connection  $A$  in  $\mathcal{A}$  defines the desired homomorphism simply through the holonomy operation:  $\hat{H}(\tilde{\alpha}) := \mathcal{P} \exp G \oint_{\tilde{\alpha}} A \cdot dl$ , where  $\alpha$  is any loop in the hoop  $\tilde{\alpha}$ . However, there are many homomorphisms which do not arise from smooth connections. This leads to “generalized connections” –i.e. elements in  $\overline{\mathcal{A}/\mathcal{G}} - \mathcal{A}/\mathcal{G}$ . In particular, there exist  $\bar{A}$  in  $\overline{\mathcal{A}/\mathcal{G}}$  which have support at a single point and are thus “distributional.” Note that this characterization of the spectrum  $\overline{\mathcal{A}/\mathcal{G}}$  is *completely algebraic*; there is no continuity assumption on the homomorphisms. This property makes the characterization very useful in practice.

As noted above, positive linear functionals on  $\mathcal{HA}$  are in 1-1 correspondence with regular measures on (the compact Hausdorff space)  $\overline{\mathcal{A}/\mathcal{G}}$ . It turns out that the positive linear functions, in turn, are determined completely by certain “generating functionals”  $\Gamma(\alpha)$  on the space  $\mathcal{L}_{x_o}$  of loops based at  $x_o$ :

*There is a 1-1 correspondence between positive linear functionals on  $\mathcal{HA}$  (and hence regular measures on  $\overline{\mathcal{A}/\mathcal{G}}$ ) and functional  $\Gamma(\alpha)$  on  $\mathcal{L}_{x_o}$  satisfying:*

- i)  $\sum_i a_i T_{\alpha_i} = 0 \Rightarrow \sum_i a_i \Gamma(\alpha_i) = 0$ ; and,
  - ii)  $\sum_{i,j} \bar{a}_i a_j (\Gamma(\alpha_i \circ \alpha_j) + \Gamma(\alpha_i \circ \alpha_j^{-1})) \geq 0$ .
- for all loops  $\alpha_i$  and complex numbers  $a_i$ .*

The first condition implies that the functional  $\Gamma$  is well-defined on hoops. Hence we could have taken it to be a functional on  $\mathcal{HG}$  from the beginning. Thus, we see that there is a nice “non-linear duality” between the spectrum  $\overline{\mathcal{A}/\mathcal{G}}$  and the hoop group  $\mathcal{HG}$ : Elements of  $\overline{\mathcal{A}/\mathcal{G}}$  are homomorphisms from  $\mathcal{HG}$  to  $SU(2)$  and regular measures on  $\overline{\mathcal{A}/\mathcal{G}}$  correspond to certain functionals on  $\mathcal{HG}$ . Finally, if one is interested in measures on  $\overline{\mathcal{A}/\mathcal{G}}$  which are invariant under the (induced) action of diffeomorphisms on  $\Sigma$ , one is led to seek functionals  $\Gamma(\alpha)$  which depend not on the individual loops  $\alpha$  but rather on the (generalized) knot class to which  $\alpha$  belongs. (The qualification “generalized” refers to the fact that here we are allowing the loops to have kinks, overlaps and self-intersections. Until recently, knot theorists considered only smoothly embedded loops.) Thus, there is an interesting –and potentially powerful– interplay between knot theory and representations of the holonomy algebra  $\mathcal{HA}$  in which the diffeomorphism group of  $\Sigma$  is unitarily implemented.

Finally, we can make the integration theory more explicit as follows. Consider a subgroup  $S_n$  of the hoop group  $\mathcal{HG}$  which is generated by  $n$  independent hoops. We can introduce the following equivalence relation on  $\overline{\mathcal{A}/\mathcal{G}}$ :  $\bar{A} \approx \bar{A}'$  if and only if their action on all elements of  $S_n$  coincides, i.e., if and only if  $\bar{A}(\tilde{\alpha}) = g^{-1} \cdot \bar{A}'(\tilde{\alpha}) \cdot g$  for all  $\tilde{\alpha} \in S_n$  and some (hoop independent)  $g \in SU(2)$ . Thus, intuitively, to be equivalent,  $\bar{A}$  and  $\bar{A}'$  have to agree at points on the set of loops in  $S_n$ ; their behavior elsewhere on  $\Sigma$  does not matter.

It turns out that the quotient space is isomorphic to  $[SU(2)]^n/Ad$ . This is the key result that enables us to make the space of integrable functions and the integration procedure itself more transparent. It says that the quotients obtained using two different sub-groups of the hoop group are isomorphic if and only if the two sub-groups are generated the same number of independent hoops. The details of the subgroups do not matter. There is a certain “universality” to the structure of the quotient.

We want to be able to integrate the generalized Wilson loop functionals  $T_{\tilde{\alpha}}(\bar{A}) := \bar{A}(\tilde{\alpha})$  on  $\overline{\mathcal{A}/\mathcal{G}}$ . Using the quotient construction given in the last paragraph, we will first present a convenient characterization of the space of all integrable functions on  $\overline{\mathcal{A}/\mathcal{G}}$ . Let us begin with a definition. A function  $f$  on  $\overline{\mathcal{A}/\mathcal{G}}$  will be said to be *cylindrical* if it is the pull-back to  $\overline{\mathcal{A}/\mathcal{G}}$  of a smooth function  $\tilde{f}$  on  $[SU(2)]^n/Ad$  for some sub-group  $S_n$  of the hoop group. These will be the integrable functions and they include the generalized Wilson loops. Next, we define integrals of these functions  $f$  on  $\overline{\mathcal{A}/\mathcal{G}}$  explicitly using the integrals of  $\tilde{f}$  on  $[SU(2)]^n/Ad$ , provided of course we equip  $[SU(2)]^n/Ad$  with suitable measures  $d\mu_n$  for each  $n$ . We can then first define a positive linear functional  $\Gamma'$  on the space of cylindrical functions  $f$  via:

$$\Gamma'(f) := \int_{[SU(2)]^n/Ad} \tilde{f} d\mu_n . \quad (3.2)$$

For the functional to be well-defined, of course, the family of measures  $d\mu_n$  on  $[SU(2)]^n/Ad$  must satisfy certain consistency conditions. It turns out that these requirements can be met. The resulting functionals  $\Gamma'(f)$  define regular measures  $d\mu'$  on  $\overline{\mathcal{A}/\mathcal{G}}$  such that  $\Gamma'(f) = \int f d\mu'$ .

A particularly natural choice (and, not surprisingly, the first to be discovered) is to let  $d\mu_n$  be simply induced on  $[SU(2)]^n/Ad$  by the Haar-measure on  $SU(2)$ . With this choice, we have the following results:

- i) The consistency conditions are satisfied; the left side of (3.2) is well-defined for all cylindrical functions  $f$  on  $\overline{\mathcal{A}/\mathcal{G}}$ ;
- ii) The generalized holonomies  $T_{\tilde{\alpha}}$  are cylindrical functionals on  $\overline{\mathcal{A}/\mathcal{G}}$  and  $\Gamma(\alpha) := \Gamma'(T_{\tilde{\alpha}})$  defined via (3.2) serves as a generating functional for a faithful, cyclic representation of the homomony  $C^*$ -algebra  $\mathcal{HA}$  which ensures that  $d\mu$  is a regular, strictly positive measure on  $\overline{\mathcal{A}/\mathcal{G}}$ ;
- iii) The measure  $d\mu$  is invariant under the induced action of the diffeomorphism group on  $\Sigma$ .

(The knot invariant defined by  $d\mu$  is a genuinely generalized one; roughly, it counts the number of self-overlaps in any given loop.)

This measure is in some ways analogous to the Gaussian measure on linear vector spaces. Both can be obtained by a “cylindrical construction.” The Gaussian measure uses the natural metric on  $\mathbb{R}^n$  while the above measure uses the natural (induced) Haar measure on  $[SU(2)]^n/Ad$ . They are both regular and strictly positive. This leads us to ask if other properties of the Gaussian measure are shared. For instance, we know that the Gaussian measure is concentrated on distributions; although the smooth fields are dense in the space of distributions in an appropriate topology, they are contained in a set whose

total measure is zero. Is the situation similar here? The answer turns out to be affirmative. The classical configuration space  $\mathcal{A}/\mathcal{G}$  with which we began is dense in the domain space  $\overline{\mathcal{A}/\mathcal{G}}$  of quantum states in the Gel'fand topology. However,  $\mathcal{A}/\mathcal{G}$  is contained in a set whose total measure is zero. The measure is again concentrated on “generalized” connections in  $\overline{\mathcal{A}/\mathcal{G}}$  (Marolf & Mourão (1993)). In a certain sense, just as the Gaussian measures on linear spaces originate in the harmonic oscillator, the new measure on  $\overline{\mathcal{A}/\mathcal{G}}$  originates in a (generalized) rotor (whose configuration space is the  $SU(2)$  group-manifold). However, the measure is, so to say, “genuinely” tailored to the underlying non-linearity. It is *not* obtained by “perturbing” the Gaussian measure.

With the measure  $d\mu$  at hand, we can consider the Hilbert space  $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu)$  and introduce operators on it. This is *not* the Hilbert space of physical states of quantum gravity since we have not imposed constraints. It is a fiducial, *kinematical* space which is to enable us to regularize various operators (in particular, the quantum constraint operators). The configuration operators are associated with the generalized Wilson loop functionals:  $\hat{T}_{\tilde{\alpha}} \circ \Psi(\bar{A}) = \bar{A}(\tilde{\alpha})\Psi(\bar{A})$ . One can show that there are bounded, self-adjoint operators on the Hilbert space. There are also “momentum operators” –associated with closed, 2-dimensional ribbons or strips in the 3-manifold  $\Sigma$ – which are gauge invariant and linear in the electric field. One can show that these are also self-adjoint (but unbounded). Finally, since  $d\mu$  is invariant under the induced action of the diffeomorphism group of  $\Sigma$ , this group acts unitarily.

Next, we can now make the Rovelli-Smolín loop transform (Rovelli & Smolin (1990), see, also Gambini & Trias (1986), Brüggmann (1994), Loll (1994)) rigorous. Since each generalized Wilson loop function  $T_{\tilde{\alpha}}(\bar{A})$  is cylindrical and belongs to the Hilbert space  $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu)$ , the loop transform can be defined simply as a scalar product:

$$\psi(\tilde{\alpha}) := \int_{\overline{\mathcal{A}/\mathcal{G}}} \overline{T_{\tilde{\alpha}}(\bar{A})} \Psi(\bar{A}) d\mu . \quad (3.3)$$

As desired, it enables us to pass from functionals  $\Psi(\bar{A})$  of generalized connections to functionals  $\psi(\tilde{\alpha})$  of hoops<sup>†</sup>. With the rigorously defined transform at hand, we can take over various operators from the connection to the loop representation. For the Wilson loop operators  $\hat{T}_{\tilde{\alpha}}^0$  as well as the strip operators –which are the smeared versions of  $\hat{T}_{\tilde{\alpha}}^a(s)$ – we recover the same answers as have been used in all the calculations in the loop representation. For the operator  $\hat{T}^{ab}(s, t)$ , which is used in the regularization of the metric and of the Hamiltonian constraint operators, work is still in progress.

Finally, let us consider the constraints. Our task here is to represent the quantum constraints as well-defined operators on the Hilbert space  $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu)$ , take them over to the loop side and solve them. For the diffeomorphism constraint, most of this task has been carried out. The operator is well-defined and self-adjoint on the connection states

---

<sup>†</sup> Since the left side is a function of hoops, we should, strictly, use the terms hoop representation and the hoop transform. However, in various calculations, it is often convenient to lift these functionals from the hoop group  $\mathcal{HG}$  to the space of loops  $\mathcal{L}_{x_o}$ . Therefore, as in most of the literature on the subject, we will not keep a careful distinction between loops and hoops in what follows.



and can be taken over to the loop side unambiguously. The result is the expected one: the operator acts on the loop states  $\psi(\alpha)$  by displacing the loop  $\alpha$  in the argument via the corresponding diffeomorphism. Its functional form is the one used in the literature. It is therefore intuitively clear that the solutions to this constraint functions of (generalized) knot classes. The question is: What type of functions are we to allow? As explained by Petr Hajiček (1994), typically, zero is a continuous eigenvalue of the constraint operators whence elements of the kernel are not normalizable. Thus, the availability of the Hilbert space by itself does not provide us with the regularity conditions that must be satisfied by the *physical* states. For this, as Hajiček pointed out, we need a rigged Hilbert space construction. To fix the ideas, let me recall the situation in the case of a relativistic particle, where the classical constraint is  $P^\alpha P_\alpha + \mu^2 = 0$ . In this case, the kinematical Hilbert space can be taken to be  $L^2(\mathbb{R}^4)$ . This space is needed to translate the classical constraint function to a well-defined operator (whose kernel can then be found). The operator, of course, is  $-\hbar^2 \eta^{\alpha\beta} \partial_\alpha \partial_\beta + \mu^2$ . It is self-adjoint on the kinematic Hilbert space  $L^2(\mathbb{R}^4)$ . However, no (non-zero) element in its kernel is normalizable. These elements belong to the rigged Hilbert space; in the momentum space, they are distributions with support on the mass shell. To find out the regularity conditions on the (generalized) knot invariants, we need to carry out a similar construction and isolate the appropriate rigged Hilbert space. Work is in progress on this problem.

For the Hamiltonian constraint, there is still no progress to report on the mathematical side. This is not surprising in the light of the fact that work on the operators  $\hat{T}^{ab}(s, t)$  is still in progress. But the general line of attack is clear. At the heuristic level, there have been several distinct approaches to the problem of constructing this operator in the loop representation but they have all led to equivalent results (Brömann & Pullin (1993)). In an optimistic view, this could be taken as a signal that the operator *would* exist rigorously.

To summarize then, there has been a considerable amount of rigorous work in the last two years and the goal of giving a precise meaning to the loop transform and the constraint operators as well as that of extracting the regularity conditions on the physical states now seems attainable in the near future. The main open problems are: i) extending the theory to incorporate complex connections; ii) writing the Hamiltonian constraint as a well-defined operator on  $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu)$ ; and, iii) constructing the appropriate rigged Hilbert spaces to obtain regularity conditions on the physical states. Work is in progress on all these problems.

Thus, it appears that the integration theory based on the measure  $d\mu$  will provide the mathematical basis for this non-perturbative approach to quantum gravity. However, these techniques may be used also in other theories of connections which are diffeomorphism invariant and perhaps even in Yang-Mills theory which is not diffeomorphism invariant. We saw that the full domain space of quantum theory,  $\overline{\mathcal{A}/\mathcal{G}}$ , can be thought of as the space of homomorphisms from the full hoop group  $\mathcal{H}\mathcal{G}$  to the gauge group  $SU(2)$ . Given a finitely generated sub-group  $S_n$  of the hoop group, we can consider the space of homomorphisms from it to  $SU(2)$ . This provides the space  $[SU(2)]^n / Ad$  which is precisely the domain space of quantum states of a lattice gauge theory where the lattice is not rectangular but tailored to the given subgroup  $S_n$  of the gauge group. Thus, what we have is a set of “floating lattices,” each associated with a finitely generated subgroup of the hoop group. The space

$\overline{\mathcal{A}/\mathcal{G}}$  can be rigorously recovered as a projective limit of the configuration spaces of lattice theories (Marolf & Mourão 1993). This construction is potentially quite powerful; it may enable one to take continuum limits of operators of lattice theories in a completely new fashion. The limit is obtained not by taking the lattice separation to zero but by enlarging lattices to probe the continuum connections better and better, i.e., by considering larger and larger subgroups of the hoop group. This is a good illustration of the synergistic exchange of ideas between general relativity and gauge theories that is made possible by this approach to quantum gravity.

#### 4. Emergence of low energy physics

One of the important features that any non-perturbative approach to quantum gravity must have is that, in a suitable limit, it should reproduce the laboratory scale physics correctly. Does the theory admit states which can be interpreted as smooth geometries on large scales? Is there at least an approximate notion of time which is compatible with a space-time picture? Can one regard non-gravitational physical fields as evolving unitarily with respect to this time? Is there an approximate Hamiltonian governing this evolution? One often takes for granted that the answers to such questions would be obviously “yes.” However, in genuinely non-perturbative treatments, this is by no means clear a priori; one may be working in a sector of the quantum theory which does not admit the correct or unambiguous classical limit. For example, the sector may correspond to a confined phase which has no classical analog or the limit may yield a wrong number even for the macroscopic dimensions of space-time!

In this section, I will present two examples of such results that have been obtained within the present non-perturbative framework which were not discussed by other speakers. The first involves the existence of quantum states which approximate classical geometries at macroscopic scales (Ashtekar et al, 1992). I will begin by showing that certain operators representing *geometrical observables* can be regulated in a way that respects the diffeomorphism invariance of the underlying theory. What is more, these regulated operators are finite *without any renormalization*. Using these operators, one can ask if there exist loop states which approximate smooth geometry at large scales. Not only is the answer in the affirmative but, furthermore, *these states exhibit a discrete structure of a definite type at the Planck scale*. (For further details, see, e.g., Rovelli & Smolin (1990), Ashtekar (1992) and Smolin (1993).) The second example involves the issue of time in a certain weak field truncation of the theory. Here, I will only summarize overall the situation, emphasizing those points which will be needed in the next section (For details, see, Ashtekar (1991), chapter 12 and, for more recent developments, Rovelli and Smolin (1993).)

##### 4.1 Non-perturbative regularization

Let us begin with the issue of regularization. As noted by Guilliani (1994), in the present framework, the spatial metric is constructed from products of “electric fields”  $E_i^a$ . It is thus a “composite” field given by  $q^{ab}(x) = E^{ai}(x)E_i^b(x)$ . (This field is, as emphasized by other speakers, a density of weight 2. To keep the notation simple, however, I will drop the tildes that are generally used to denote the density weights.) In the quantum theory, therefore, this operator must be regulated. The obvious possibility is point splitting. One might set  $q^{ab}(x) = \lim_{y \rightarrow x} E^{ai}(x)E_i^b(y)$ . However, the procedure violates gauge invariance

since the internal indices at two *different* points have been contracted. There is, however, a suitable modification that will ensure gauge invariance. Consider the field  $T^{aa'}[\alpha](y', y)$ , labelled by a closed loop  $\alpha$  and points  $y$  and  $y'$  thereon, defined in the classical theory by:

$$T^{aa'}[\alpha](y, y') := \frac{1}{2} \text{Tr} \left[ (\mathcal{P} \exp G \int_{y'}^y A_b dl^b) E^a(y') (\mathcal{P} \exp G \int_y^{y'} A_c dl^c) E^{a'}(y) \right]. \quad (4.1)$$

In the limit  $\alpha$  shrinks to zero,  $T^{aa'}[\alpha](y, y')$  tends to  $-4q^{aa'}$ .

Now, in quantum theory, one can define the action of the operator  $\hat{T}^{aa'}[\alpha](y, y')$  directly on the loop states  $\psi(\beta)$ . The explicit form will not be needed here. We only note that using the bra-ket notation,  $\psi(\beta) = \langle \beta | \psi \rangle$  the action can be specified easily. Indeed,  $\langle \beta | \circ \hat{T}^{aa'}[\alpha](y, y')$  is rather simple: if a loop  $\beta$  does not intersect  $\alpha$  at  $y$  or  $y'$ , the operator simply annihilates the bra  $\langle \beta |$  while if an intersection does occur, it breaks and re-routes the loop  $\beta$ , each routing being assigned a specific weight. One may therefore try to define a quantum operator  $\hat{q}^{aa'}$  as a limit of  $\hat{T}^{aa'}[\alpha]$  as  $\alpha$  shrinks to zero. The resulting operator does exist after suitable regularization *and* renormalization. However (because of the density weights involved) the operator necessarily carries a memory of the background metric used in regularization. Thus, the idea of defining the metric operator again fails. In fact one can give general qualitative arguments to say that there are no *local*, operators which carry the metric information *and* which are independent of background fields (used in the regularization). Thus, in quantum theory, the absence of background fields introduces new difficulties. That such difficulties would arise was recognized quite early by Chris Isham and John Klauder.

There do exist, however, *non-local* operators which can be regulated in a way that respects diffeomorphism invariance.

As the first example, consider the function  $Q(\omega)$  –representing the smeared 3-metric– on the classical phase space, defined by

$$Q(\omega) := \int d^3x \ (q^{ab} \omega_a \omega_b)^{\frac{1}{2}}, \quad (4.2)$$

where  $\omega_a$  is any smooth 1-form of compact support. (Note that the integral is well-defined without the need of a background volume element because  $q^{ab}$  is a density of weight 2.) It is important to emphasize that, in spite of the notation,  $Q(\omega)$  is *not* obtained by smearing a *distribution* with a test field; because of the square-root,  $Q(\omega)$  is *not* linear in  $\omega$ . We can, nonetheless define the corresponding quantum operator as follows. Let us choose on  $\Sigma$  test fields  $f_\epsilon(x, y)$  (which are densities of weight one in  $x$  and) which satisfy:

$$\lim_{\epsilon \rightarrow 0} \int_{\Sigma} d^3x f_\epsilon(x, y) g(x) = g(y) \quad (4.3)$$

for all smooth functions of compact support  $g(x)$ . If  $\Sigma$  is topologically  $\mathbb{R}^3$ , for example, we can construct these test fields as follows:

$$f_\epsilon(x, y) = \frac{\sqrt{h(x)}}{\pi^{\frac{3}{2}} \epsilon^3} \exp - \frac{|\vec{x} - \vec{y}|^2}{2\epsilon^2}, \quad (4.4)$$

where  $\vec{x}$  are the Cartesian coordinates labeling the point  $x$  and  $h(x)$  is a “background” scalar density of weight 2. Next, let us define

$$q_\epsilon^{aa'}(x) = -\frac{1}{4} \int_\Sigma d^3y \int_\Sigma d^3y' f_\epsilon(x, y) f_\epsilon(x, y') T^{aa'}(y, y'). \quad (4.5)$$

As  $\epsilon$  tends to zero, the right side tends to  $q^{ab}$  because the test fields force both the points  $y$  and  $y'$  to approach  $x$ , and hence the loop passing through  $y, y'$ , used in the definition of  $T^{aa'}(y, y')$ , to zero. It is now tempting to try to define a local metric operator  $\hat{q}^{aa'}$  corresponding to  $q^{aa'}$  by replacing  $T^{aa'}(y, y')$  in (4.5) by its quantum analog and then taking the limit. One finds that the limit does exist provided we first renormalize  $\hat{q}_\epsilon^{aa'}$  by an appropriate power of  $\epsilon$ . However, as before, the answer depends on the background structure (such as the density  $h(x)$ ) used to construct the test fields  $f_\epsilon(x, y)$ . If, however, one tries to construct the quantum analog of the *non-local* classical variable  $Q(\omega)$ , this problem disappears. To see this, let us first express  $Q(\omega)$  using (4.5) as:

$$Q(\omega) = \lim_{\epsilon \rightarrow 0} \int_\Sigma d^3x (q_\epsilon^{aa'} \omega_a \omega_{a'})^{\frac{1}{2}}. \quad (4.6)$$

The required quantum operator  $\hat{Q}(\omega)$  on the loop states can now be obtained by replacing  $T^{aa'}(y, y')$  by the operator  $\hat{T}^{aa'}(y, y')$ . A careful calculation shows that: i) the resulting operator is well-defined on loop states; ii) no renormalization is necessary, i.e., the limit is automatically *finite*; and, iii) the final answer carries no imprint of the background structure (such as the density  $h(x)$  or, more generally, the specific choice of the test fields  $f_\epsilon(x, y)$ ) used in regularization. To write out its explicit expression, let me restrict myself to smooth loops  $\alpha$  without any self-intersection. Then, the action is given simply by:

$$\langle \alpha | \circ \hat{Q}(\omega) = l_P^2 \oint_\alpha ds |\dot{\alpha}^a \omega_a| \cdot \langle \alpha |, \quad (4.7)$$

where  $l_P = \sqrt{G\hbar}$  is the Planck length,  $s$ , a parameter along the loop and  $\dot{\alpha}^a$  the tangent vector to the loop. In this calculation, the operation of taking the square-root is straightforward because the relevant operators are diagonal in the loop representation. This is analogous to the fact that, in the position representation of non-relativistic quantum mechanics, we can set  $\langle x | \circ \sqrt{\hat{X}^2} = \langle x | \cdot |x|$  without recourse to the detailed spectral theory. The  $G$  in  $l_P$  of (4.7) comes from the fact that  $GA_a^i$  has the usual dimensions of a connection while  $\hbar$  comes from the fact that  $\hat{E}_i^a$  is  $\hbar$  times a functional derivative. The final result is that, on non-intersecting loops, the operator acts simply by multiplication: the loop representation is well-suited to find states in which the 3-geometry –rather than its time evolution– is sharp.

The second class of operators corresponds to the area of 2-surfaces. Note first that, given a smooth 2-surface  $S$  in  $\Sigma$ , its area  $\mathcal{A}_S$  is a function on the classical phase space. We first express it using the classical loop variables. Let us divide the surface  $S$  into a large number  $N$  of area elements  $S_I, I = 1, 2, \dots, N$ , and set  $\mathcal{A}_I^{\text{appr}}$  to be

$$\mathcal{A}_I^{\text{appr}} = -\frac{1}{4} \left[ \int_{S_I} d^2S^{bc}(x) \eta_{abc} \int_{S_I} d^2S^{b'c'}(x') \eta_{a'b'c'} T^{aa'}(x, x') \right]^{\frac{1}{2}}, \quad (4.8)$$

where  $\eta_{abc}$  is the (metric independent) Levi-Civita density of weight  $-1$ . It is easy to show that  $\mathcal{A}_I^{\text{appr}}$  approximates the area function (on the phase space) defined by the surface elements  $S_I$ , the approximation becoming better as  $S_I$  –and hence loops with points  $x$  and  $x'$  used in the definition of  $T^{aa'}$ – shrink. Therefore, the total area  $\mathcal{A}_S$  associated with  $S$  is given by

$$\mathcal{A}_S = \lim_{N \rightarrow \infty} \sum_{I=1}^N \mathcal{A}_I^{\text{appr}}. \quad (4.9)$$

To obtain the quantum operator  $\hat{\mathcal{A}}_S$ , we simply replace  $T^{aa'}$  in (4.8) by the quantum loop operator  $\hat{T}^{aa'}$ . This somewhat indirect procedure is necessary because, as indicated above, there is no well-defined operator-valued distribution that represents the metric or its area element *at a point*. Again, the operator  $\hat{\mathcal{A}}_S$  turns out to be finite. Its action, evaluated on a nonintersecting loop  $\alpha$  (for simplicity), is given by:

$$\langle \alpha | \circ \hat{\mathcal{A}}_S = \frac{l_p^2}{2} I(S, \alpha) \cdot \langle \alpha |, \quad (4.10)$$

where  $I(S, \alpha)$  is simply the *unoriented* intersection number between the 2-surface  $S$  and the loop  $\alpha$ . (One obtains the *unoriented* intersection number here and the absolute sign in the integrand of (4.7) because of the square-root operation involved in the definition of these operators.) Thus, in essence, “a loop  $\alpha$  contributes half a Planck unit of area to any surface it intersects.”

The fact that the area operator also acts simply by multiplication on non-intersecting loops lends further support to the idea that the loop representation is well-suited to “diagonalize” operators describing the 3-geometry. Indeed, we can immediately construct a large set of simultaneous eigenbras of the smeared metric and the area operators. There is one,  $\langle \alpha |$ , associated to every nonintersecting loop  $\alpha$ . Note that the corresponding eigenvalues of area are *quantized* in integral multiples of  $l_p^2/2$ . There are also eigenstates associated with intersecting loops which, however, I will not go into to since the discussion quickly becomes rather involved technically.

## 4.2 Weaves

With these operators on hand, we can now turn to the construction of quantum loop states that approximate the classical metric  $h_{ab}$  on  $\Sigma$  on a scale large compared to the Planck length. The basic idea is to weave the classical metric out of quantum loops by spacing them so that, on an average, precisely one line crosses *any* surface element whose area, *as measured by the given  $h_{ab}$*  is one Planck unit. Such loop states will be called *weaves*. Note that these states are not uniquely picked out since our requirement is rather weak. Indeed, given a weave approximating a specific classical metric, one can obtain others, approximating the same classical metric.

Let us begin with a concrete example of such a state which will approximate a *flat* metric  $h_{ab}$ . To construct this state, we proceed as follows. Using this metric, let us introduce a random distribution of points on  $\Sigma = \mathbb{R}^3$  with density  $n$  (so that in any given volume  $V$  there are  $nV(1 + \mathcal{O}(1/\sqrt{nV}))$  points). Center a circle of radius  $a = (1/n)^{1/3}$  at each of these points, with a random orientation. We assume that  $a \ll L$ , so that there is

a large number of (non-intersecting but, generically, *linked*) loops in a macroscopic volume  $L^3$ . Denote the collection of these circles by  $\Delta_a$ . As noted in section 4, due to trace identities, products of Wilson loop functionals  $T_{\tilde{\alpha}}$  can be expressed as linear combinations of Wilson loop functionals. As a consequence, it turns out that the bras defined by multi-loops are equivalent to linear combinations of single loop bras. Therefore, for each choice of the parameter  $a$ , there is a well-defined bra  $\langle \Delta_a |$ . This is our candidate weave state.

Let us consider the observable  $\hat{Q}[\omega]$ . To see if  $\langle \Delta_a |$  reproduces the geometry determined by the classical metric  $h_{ab}$  on a scale  $L \gg l_p$ , let us introduce a 1-form  $\omega_a$  which is *slowly varying on the scale  $L$*  and compare the value  $Q[\omega](h)$  of the classical  $Q[\omega]$  evaluated at the metric  $h_{ab}$ , with the action of the quantum operator  $\hat{Q}[\omega]$  on  $\langle \Delta_a |$ . A detailed calculation yields:

$$\langle \Delta_a | \circ \hat{Q}[\omega] = \left[ \frac{\pi}{2} \left( \frac{l_p}{a} \right)^2 Q[\omega](h) + \mathcal{O}\left(\frac{a}{L}\right) \right] \cdot \langle \Delta_a |. \quad (4.11)$$

Thus,  $\langle \Delta_a |$  is an eigenstate of  $\hat{Q}[\omega]$  and the corresponding eigenvalue is closely related to  $Q[\omega](h)$ . However, even to the leading order, the two are unequal *unless* the parameter  $a$  –the average distance between the centers of loops– *equals*  $\sqrt{\pi/2} l_p$ . More precisely, (4.11) can be interpreted as follows. Let us write the leading coefficient on the right side of this equation as  $(1/4)(2\pi a/l_p)(nl_p^3)$ . Since this has to be unity for the weave to reproduce the classical value (to leading order), we see that  $\Delta_a$  should contain, on an average, one fourth Planck length of curve per Planck volume, where lengths and volumes are measured using  $h_{ab}$ .

The situation is the same for the area operators  $\hat{\mathcal{A}}_S$ . Let  $S$  be a 2-surface whose extrinsic curvature varies slowly on a scale  $L \gg l_p$ . One can evaluate the action of the area operator on  $\langle \Delta_a |$  and compare the eigenvalue obtained with the value of the area assigned to  $S$  by the given flat metric  $h_{ab}$ . Again, the eigenvalue can be re-expressed as a sum of two terms, the leading term which has the desired form, except for an overall coefficient which depends on the mean separation  $a$  of loops constituting  $\Delta_a$ , and a correction term which is of the order of  $\mathcal{O}(\frac{a}{L})$ . We require that the coefficient be so adjusted that the leading term agrees with the classical result. This occurs, again, precisely when  $a = \sqrt{\pi/2} l_p$ . It is interesting to note that the details of the calculations which enable one to express the eigenvalues in terms of the mean separation are rather different for the two observables. In spite of this, the final constraint on the mean separation is *precisely* the same.

Let us explore the meaning and implications of these results.

- 1) It is generally accepted that, to obtain classical behavior from quantum theory, one needs two things: i) an appropriate coarse graining, and, ii) special states. In our procedure, the slowly varying test fields  $\omega_a$  and surfaces  $S$  with slowly varying extrinsic curvature enable us to perform the appropriate coarse graining while weaves –with the precisely tuned mean separation  $a$ – are the special states. There is, however, something rather startling: The restriction on the mean separation  $a$  –i.e., on the *short distance* behavior of the multi-loop  $\Delta_a$ – came from the requirement that  $\langle \Delta_a |$  should approximate the classical metric  $h_{ab}$  on *large scales  $L$* !
- 2) In the limit  $a \rightarrow \infty$ , the eigenvalues of the two operators on  $\langle \Delta_a |$  go to zero. This is not too surprising. Roughly, in a state represented by any loop  $\alpha$ , one expects the

quantum geometry to be excited just at the points of the loops. If the loops are *very* far away from each other as measured by the fiducial  $h_{ab}$ , there would be macroscopic regions devoid of excitations where the quantum geometry would seem to correspond to a zero metric.

- 3) The result of the opposite limit, however, *is* surprising. One might have naively expected that the best approximation to the classical metric would occur in the continuum limit in which the separation  $a$  between loops goes to zero. However, the explicit calculation outlined above shows that this is not the case: as  $a$  tends to zero, the leading terms in the eigenvalues of  $\hat{Q}[\omega]$  and  $\mathcal{A}_S$  actually diverge! (One's first impulse from lattice gauge theories may be to say that the limit is divergent simply because we are not rescaling, i.e., renormalizing the operator appropriately. Note, however, that, in contrast to the calculations one performs in lattice theories, here, we *already* have a well defined operator in the continuum. We are only probing the properties of its eigenvectors and eigenvalues, whence there is nothing to renormalize.) It is, however, easy to see the reason underlying this behavior. Intuitively, the factors of the Planck length in (4.7) and (4.10) force each loop in the weave to contribute a Planck unit to the eigenvalue of the two geometrical observables. In the limit  $a \rightarrow 0$ , the number of loops in any fixed volume (relative to the fiducial  $h_{ab}$ ) grows unboundedly and the eigenvalue diverges.
- 4) It is important to note the structure of the argument. In non-perturbative quantum gravity, there is no background space-time. Hence, terms such as “slowly varying” or “microscopic” or “macroscopic” have, a priori, no physical meaning. One must do some extra work, introduce some extra structure to make them meaningful. The required structure should come from the very questions one wants to ask. Here, the questions had to do with approximating a classical geometry. Therefore, we could *begin* with classical metric  $h_{ab}$ . We used it repeatedly in the construction: to introduce the length scale  $L$ , to speak of “slowly varying” fields  $\omega_a$  and surfaces  $S$ , and, to construct the weave itself. The final result is then a consistency argument: If we construct the weave according to the given prescription, then we find that it approximates  $h_{ab}$  on macroscopic scales  $L$  provided we choose the mean separation  $a$  to be  $\sqrt{\pi/2}l_P$ , where all lengths are measured relative to the same  $h_{ab}$ .
- 5) Note that there is a considerable non-uniqueness in the construction. As we noted already, a given 3-geometry can lead to distinct weave states; our construction only serves to make the existence of such states explicit. For example, there is no reason to fix the radius  $r$  of the individual loops to be  $a$ . For the calculation to work, we only need to ensure that the loops are large enough so that they are generically linked and small enough so that the values of the slowly varying fields on each loop can be regarded as constants plus error terms which we can afford to keep in the final expression. Thus, it is easy to obtain a 2-parameter family of weave states, parametrized by  $r$  and  $a$ . The condition that the leading order terms reproduce the classical values determined by  $h_{ab}$  then gives a relation between  $r$ ,  $a$  and  $l_P$  which again implies discreteness. Clearly, one can further enlarge this freedom considerably: There are a lot of eigenbras of the smeared-metric and the area operators whose eigenvalues approximate the classical values determined by  $h_{ab}$  up to terms of the

order  $\mathcal{O}(\frac{l_p}{L})$  since this approximation ignores Planck scale quantum fluctuations.

- 6) Finally, I would like to emphasize that, at a conceptual level, the important point is that the eigenvalues of  $\hat{Q}[\omega]$  and  $\mathcal{A}[S]$  can be *discrete*.

Let me conclude the discussion on weaves with two remarks. First, it is not difficult to extend the above construction to obtain weave states for curved metrics  $g_{ab}$  which are slowly varying with respect to a flat metric  $h_{ab}$ . Given such a metric, one can find a slowly varying tensor field  $t_a{}^b$ , such that the metric  $g_{ab}$  can be expressed as  $t_a{}^c t_b{}^d h_{cd}$ . Then, given a weave of the type  $\langle \Delta |$  considered above approximating  $h_{ab}$ , we can “deform” each circle in the multi-loop  $\Delta$  using  $t_a{}^b$  to obtain a new weave  $\langle \Delta |_t$  which approximates  $g_{ab}$  in the same sense as  $\langle \Delta |$  approximates  $h_{ab}$ . (See, also, Zegwaard (1993, 1994) for the weave corresponding to the Schwarzschild black-hole.) The second remark is that since the weaves are eigenbras of the operators that capture the 3-geometry, they do not approximate 4-geometries. To obtain a state that can approximate Minkowski space-time, for example, one has to consider a loop state that resembles a “coherent state” peaked at the weave  $\Delta_a$ . In that state, neither the 3-geometry nor the time-derivative thereof would be sharp; but they would have minimum spreads allowed by the uncertainty principle. This issue has been examined in detail by Iwasaki and Rovelli (1993).

Since these results are both unexpected and interesting, it is important to probe their origin. We see no analogous results in familiar theories. For example, the eigenvalues of the fluxes of electric or magnetic fields are not quantized in QED nor do the linearized analogs of our geometric operators admit discrete eigenvalues in spin-2 gravity. Why then did we find qualitatively different results? The technical answer is simply that the familiar results refer to the *Fock representation* for photons and gravitons while we are using a completely different representation here. Thus, the results *are* tied to our specific choice of the representation. Why do we not use Fock or Fock-like states? It is not because we insist on working with loops rather than space-time fields such as connections. Indeed, one *can* translate the Fock representation of gravitons and photons to the loop picture. (See, e.g., Ashtekar et al (1991) and Ashtekar & Rovelli, (1992).) And then, as in the Fock space, the discrete structures of the type we found in this section simply disappear. However, to construct these loop representations, one must use a flat background metric and essentially every step in the construction violates diffeomorphism invariance. Indeed, there is simply no way to construct “familiar, Fock-like” representations without spoiling the diffeomorphism invariance. Thus, the results we found are, in a sense, a direct consequence of our desire to carry out a genuinely non-perturbative quantization without introducing any background structure. However, we do not have a uniqueness theorem singling out the measure  $d\mu$  which was used to define the loop transform and hence to construct the loop representation used here. There do exist other diffeomorphism invariant measures which will lead to other loop representations. The measure we have used is the most natural and the simplest among the known ones. Whether the results presented here depend sensitively on the choice of the measure is not known. Therefore, it would be highly desirable to have a uniqueness theorem which tells us that, on physical grounds, we should restrict ourselves to a specific (class of) measure(s).

My overall viewpoint is that one should simultaneously proceed along two lines: i) one should take these results as an indication that we are on the right track and push heuristic



calculations within this general framework; and, ii) one should try to put the available heuristic results on rigorous mathematical footing to avoid the danger of “wandering off” in unsound directions.

#### 4.3 The issue of time

The results reported in the last two sub-sections were kinematical; constraints were not yet imposed. This is because our questions themselves referred to 3-metrics which fail to be Dirac observables; in the classical analysis, 3-metrics are well-defined only on the full (and hence kinematical) phase space rather than on the reduced phase space (the space of physical or “true” degrees of freedom). On the other hand, as Hajiček (1994) explained in his lectures, the issue of time is *not* kinematical; it is intimately intertwined with the scalar or the Hamiltonian constraint. Therefore, we must now bring in this constraint. We will show that in a certain truncation of the full quantum theory, the issue of time can be resolved satisfactorily in the asymptotically flat context.

It turns out that, for technical reasons sketched below, (in pure gravity) it is the connection representation that seems best-suited to handle this issue. Thus, the question we wish to now ask is the following: can we single out a component of the connection,  ${}^TA$  which can serve as “internal” time? More precisely, can we re-interpret the scalar –or Hamiltonian– constraint as telling us how the “true” or the dynamical degrees of freedom  ${}^DA$  evolve with respect to the internal clock  ${}^TA$ ? The questions themselves are not new. They have been asked in the context of geometrodynamics since the sixties. It turned out, however, that one can not isolate time in this manner in the metric representation of geometrodynamics even in the weak field truncation (Kuchař (1970)). Why does one then hope that the situation would be any better in connection dynamics?

Let us begin by analyzing the source of the difficulty. Consider, first, a parametrized, non-relativistic system. Let the phase space coordinates be  $(q^1, \dots, q^n; p_1, \dots, p_n)$  and let the constraint be:

$$p_1 + H(q^1, q^2, \dots, q^n; p_2, \dots, p_n) = 0. \quad (4.12)$$

Now, if in the quantum theory, we used the  $q$ -representation, the quantum constraint has the same form as the time-dependent Schrödinger equation,  $i\hbar\partial\Psi(q^i)/\partial q^1 = \hat{H} \circ \Psi(q^i)$ , where  $q^1$  plays the role of time. On the other hand, the momentum representation is equally viable mathematically. The quantum constraint would now take the form  $\hat{p}^1\Psi(p^i) = -\hat{H} \circ \Psi(p^i)$ ; it does *not* resemble the time-dependent Schrödinger equation. In particular, it is now hard to isolate time from among the configuration variables on which the wave function depends. Thus, although two representation may be mathematically equivalent, one may be better suited to deal with the issue of time. Since the Dirac transformation theory enable us to pass from one to another, the issue here is that of convenience. However, since so many of the issues related to the interpretation of the framework hinge on the availability of a (possibly approximate) time variable, a representation which is well-suited to extract an internal clock has central importance. Let us now return to gravity. Recall from Beig’s lectures that, in the asymptotically flat case, the Hamiltonian of geometrodynamics can be written as  $H_N(q, p) = C_N(q, p) + \mathcal{E}(q)$ , where  $N$  is any lapse which tends to 1 at infinity at an appropriate rate and  $\mathcal{E}(q)$  is the ADM energy integral which depends only on the 3-metric  $q_{ab}$  but not on the conjugate momentum  $p^{ab}$ . Hence, the scalar constraint

corresponding to this lapse can be written in the form:

$$\mathcal{E}(q) - H_N(q, p) = 0, \quad (4.13)$$

which is identical in form to (4.12), with  $-\mathcal{E}(q)$  playing the role of  $p_1$ . The similarity suggests that it is the variable that is “conjugate” to  $-\mathcal{E}(q)$  that should play the role of time. This idea is attractive also because  $\mathcal{E}(q)$  has the interpretation of the total energy of the system. Furthermore, the structure of (4.13) is unaffected by the presence of matter sources; it is universal. However, the similarity also suggests that the metric representation is not likely to be well-suited to extract time from among the gravitational degrees of freedom. The connection representation would be better suited since  $\mathcal{E}(q)$  would be a differential operator in the  $A$ -representation. We will see below that the expectation is borne out in the weak field truncation of the theory. Note finally that this argument also explains why it is unlikely that the gravitational loop variables would be well-suited to extract time: as we saw in the last two sub-sections, the loop representation is diagonal in the operators carrying information of the 3-geometry<sup>†</sup>. With these general remarks out of the way, we can now explain the precise sense in which the scalar constraint equation can be interpreted as a time dependent Schrödinger equation.

We will now assume that the underlying 2-manifold  $\Sigma$  is topologically  $\mathbb{R}^3$ . Let us begin by introducing a background point in the phase space,  $({}^oA_a^i = 0, {}^oE_i^a)$ , where  ${}^oE_i^a$  is a flat triad and expand out the fields  $\hat{A}_a^i$  and  $\hat{E}_i^a$  that appear in the constraint operators, in the powers of the deviations  $\hat{A}_a^i - 0$  and  $\hat{E}_i^a - {}^oE_i^a$ . Since we now have access to a background triad,  ${}^oE_i^a$ , it will be convenient to convert the internal indices to vector indices on all dynamical fields. Thus, quantum states can now be regarded as functionals of  $A_{ab} := A_a^i {}^oE_{ib}$ . Next, using the flat metric  ${}^oq_{ab}$  obtained from  ${}^oE_i^a$ , we can decompose the symmetric part of  $A_{ab}$  into its transverse-traceless, longitudinal and trace parts. Let us now impose the quantum constraints order by order in the deviation. The first order equations imply that the wave functions must depend only on the symmetric, transverse-traceless parts. That is, all other parts of  $\hat{A}_{ab}$  are at least of second order. Next, imposing the scalar constraint to the second order yields:

$$-\frac{2}{G} (\Delta {}^T\hat{E}) \circ \Psi({}^DA, {}^TA) = G ({}^{TT}\hat{A}_{ab})^* ({}^{TT}\hat{A}^{ab}) \cdot \Psi({}^DA, {}^TA), \quad (4.14)$$

where,  $\Delta$  is the Laplacian with respect to  ${}^oq_{ab}$ ,  ${}^T\hat{E}$  is the trace part of  $\hat{E}^{ab}$ ,  ${}^{TT}\hat{A}_{ab}$  is the (symmetric) transverse traceless part of  $\hat{A}_{ab}$ , and where, as before,  ${}^TA$  and  ${}^DA$  are the trace and the dynamical (i.e. all but trace) parts of  $A_{ab}$ . It is now natural to introduce the variable  $\tau(x)$  which is conjugate to the operator on the left side of this equation:  $\tau(x) = (-G/2)\Delta^{-1} \cdot {}^TA(x)$ , to write the equation more explicitly as:

$$\hbar \frac{\delta \Psi({}^DA, {}^TA)}{\delta \tau(x)} = G ({}^{TT}\hat{A}_{ab})^* ({}^{TT}\hat{A}^{ab}) \cdot \Psi({}^DA, {}^TA). \quad (4.15)$$

---

<sup>†</sup> If we couple gravity to matter, however, the picture can change. It is then possible that one of the matter variables can play the role of an internal clock. This is an attractive strategy especially in the spatially compact case.

Eq (4.15) tells us how to determine the value of the physical state  $\Psi(^DA, ^TA)$  *everywhere* on the configuration space from its value on one  $^TA = \text{const.}$  hypersurface. In this sense, the equation can be thought of as a quantum constraint. It tells us that the physical states are not freely specifiable; their functional form is constrained. To obtain the “evolution” interpretation, it suffices to use just “one component” of this functional differential equation. The idea is simply to integrate the equation on  $\Sigma$  which corresponds to taking its moment with a lapse  $N = 1$ . We can simplify the left side of the resulting equation using the fact that the states are all *holomorphic* functionals of  $A_{ab}$  and, the right side, by going to the momentum space. The result is:

$$i\hbar \left( \int d^3x \frac{\delta}{\delta \text{Im}\tau(x)} \right) \circ \Psi(^DA, ^TA) = \hbar \left( \int d^3k |k| \text{}^{TT}A_{ab}(k) \frac{\delta}{\delta \text{}^{TT}A_{ab}(k)} \right) \circ \Psi(^DA, ^TA). \quad (4.16)$$

Thus, the imaginary part,  $\text{Im}\tau(x)$ , of  $\tau(x)$  plays the role of time. The operator on the right side is precisely the Hamiltonian of the truncated theory. Thus, in the second order truncation, the scalar constraint reproduces the time-dependent Schrödinger equation. This feature remains intact if one includes matter fields as sources.

As I have emphasized, in non-perturbative quantum gravity, we do not have access to a classical space-time. Even in the above truncation procedure, we worked only with the 3-manifold  $\Sigma$  and expanded operators around some background fields on  $\Sigma$ . Thus, we did not have access to a classically defined time variable. Rather, we were able to isolate, from among the mathematical variables contained in  $A_a^i$ , a preferred time  $t$  which serves as an “internal clock” with respect to which the wave function evolves. Put differently, by identifying time in the components of  $A_a^i$ , we have derived the Schrödinger equation of weak-field gravity without having a direct access to a space-time metric or even a 4-manifold. After having obtained the result, we can look back and see that we could have obtained the same result in a space-time picture. That is, the true dynamics takes place in the infinite dimensional configuration space of connections. However, *in suitable approximations* that dynamics can be re-interpreted as taking place in a suitably constructed 4-dimensional space-time. There may, however, be instances when the approximation would break down and no space-time picture is adequate. However, if any non-perturbative description is to be viable at all, it better be the case that it reduces to Minkowskian physics with Schrödinger evolution for quantum fields in an appropriate limit. As Kuchař (1970) pointed out almost 25 years ago, this is hard to achieve directly in geometrodynamics. Connection dynamics, as we have seen, is better suited for this task.

Finally, there is another approach to the extraction of time: coupling to matter fields. Recently, Rovelli and Smolin (1993) have made significant progress by coupling gravity to a scalar and other matter fields and using the scalar field to define the internal clock. The remarkable aspect of the development is that the resulting *true* quantum Hamiltonian is a manageable operator in the loop representation. The precise domain in which time can be so extracted –and hence, the nature of the implicit approximation scheme– is, however, not clear yet. Nonetheless, the fact that concrete calculations can be done in the full, non-truncated theory, is already quite exciting. The problem of time is an old one and much effort has gone into trying to find a clean, exact solution to that problem. These approximation methods and concrete calculations may well provide the type of insight that has been missing.

## 5. Outlook

As articles in this volume illustrate, in recent years, there has been notable progress on some of the difficult issues in quantum gravity. In particular, we have seen that connection dynamics offers a unified mathematical framework for all four fundamental interactions. Not only do the loop space methods enable us to go back and forth successfully between gauge theories and gravity but they have also made contact with to the fertile area of knot theory. On the mathematical side, the subject has achieved a surprising degree of maturity over the last two years and one can look forward to further contributions from the mathematical physics community –the constructive field theorists, the  $C^*$ -algebra community, topologists and knot theorists– in the coming years.

As Brügmann discussed in some detail, the program is of course far from being complete. But the situation in 2+1-dimensional gravity suggests that the various difficulties we face may be primarily technical. With enough work, in the near future we should be able to establish that either the diverse goals of the program can in fact be met or that the program encounters *specific* unsurmountable obstacles. The second type of result would also represent concrete progress. After all, the program is driven by a “radical conservatism” —it is based on the well established principles of general relativity and quantum theory and does not begin by guessing what the Plank scale structure ought to be. Therefore, if it runs into specific problems, it is likely that these limitations themselves will suggest the required modifications.

What directions is the program likely to take in the near future? It is perhaps of interest to list a few open problems whose solution will considerably improve the current state of the field. This is *not* an exhaustive list. My primary goal is to illustrate, with concrete examples, promising directions in the hope is that these examples will stimulate young researchers as well as experts in various areas to contribute to the field:

- 1) *Mathematical issues:* As we saw in section 3, the level of mathematical precision in the Rovelli-Smolín loop transform has improved considerably. However, very little is known regarding the inverse transform. Can we characterize the loop functionals in the image of the transform in a simple way without having to refer to the connection representation? Can we define a measure on the hoop group and define the inverse transform? Does the equivalent of the Plancherel theorem –which makes the Fourier transform so powerful– hold? A promising direction is being pursued by Gambini and his collaborators where the hoops group is replaced by a group of “smoothened out” hoops (Di Bortolo et al, 1993). This group has the structure of a Lie group and offers a new approach to the problem of regularization of various operators. This may well be the framework needed to make the inverse transform well-defined.
- 2) *Euclideanization:* In constructive quantum field theory, the developments which culminated in the Osterwalder-Schrader system of Euclidean axioms caused a burst of activity. Suddenly, a variety of powerful techniques became available. The key result was the demonstration of the equivalence of the Osterwalder-Schrader system to the Wightman system: if we have a theory satisfying the first set, there exists a theory satisfying the second, even though it may be hard in practice to construct it explicitly. Thus, it is still the Lorentzian theory that is of physical interest; it is just that the mathematical problem of constructing such a theory can be reduced to a more

tractable problem in the Euclidean framework. Does an analogous Euclidean system exist for general relativity? We know that naive attempts at trying to get at the physical, Lorentzian theory via the obvious Euclidean constructions can not succeed (Mena-Marugan, 1993). Are there more subtle constructions? The discovery of such a procedure would be a key contribution since many of the technically hard problems simplify enormously in the Euclidean domain.

- 3) *Midisuperspaces*: The midi-superspace of solutions to (3+1-dimensional) Einstein's equation with one (space-like) Killing field is an especially fertile area for future work. Mathematically, this system is equivalent to 2+1-dimensional general relativity coupled to a non-linear sigma-model. If the Killing field in question is hypersurface orthogonal, the matter field reduces to a single scalar field satisfying the wave equation. (If, in addition, the norm of the Killing field is constant, the matter field disappears and we have vacuum 2+1 gravity, which we already know how to quantize.) Thus, in these midi-superspaces, the problem reduces to that of quantizing 2+1 gravity with matter. It is a genuine field theory with an infinite number of degrees of freedom. However, it has two key simplifications. First, the reality conditions now become significantly simpler: through an "internal Wick-rotation" along the Killing direction, one is led to consider *real*  $SU(1,1)$  connections in place of the complex  $SU(2)$  ones (Ashtekar & Varadarajan, 1992a). Second, in the asymptotically flat case, the total Hamiltonian of the system is (not only non-negative but also) *bounded from above* indicating strongly that the quantum theory may be finite (Ashtekar & Varadarajan, 1992b). This midi-superspace thus has just the right blend of technical simplifications and physical generality to serve as a valuable stepping stone to the full theory.
- 4) *Solutions to the Hamiltonian constraint*: The "smoothened loop" techniques introduced by Gambini and his collaborators have led to interesting new solutions to the Hamiltonian constraints which are related to some well-known knot invariants (Brügmann et al, 1992a,b). Furthermore, it is clear that these are just the simplest applications of the framework. These methods are powerful and have opened up a new line of attack to the problem of solving the Hamiltonian constraint. Obtaining explicit solutions and understanding their physical meaning is important because that would strengthen our intuition considerably. However, it is also highly desirable to get a handle on the *structure* of the space of solutions. To define the inner-product on physical states, for example, it is not necessary to have the solutions explicitly. What we need is an understanding of the various mathematical structures that naturally exist on this space. So far, this issue has attracted very little attention. Finally, recently Barbero-Gonzalez (1993) has recast general relativity in 3 and 4 dimensions as a theory of *two* connections in which the triad field is secondary and does not even enter the basic equations. This is an exciting development with a lot of potential and may, in particular, open up new avenues to the issue of solving the quantum constraints.
- 5) *Approximation methods* To address physical issues, we need to develop the approximation methods further. For example, we saw in section 4.3 that by a truncation procedure around a Minkowskian initial data, one can recover the familiar laboratory physics from the scalar constraint. An obvious question then is what would happen if we truncated around a black-hole initial data. Can one automatically recover the

Hawking effect? There are indications that one would. The analysis of “embroidery around weaves” due to Iwasaki and Rovelli (1993) is likely to play an important role in this procedure. The final result will be a derivation of the Hawking effect from the loop representation of full quantum gravity (i.e., from “above”) rather than from quantum field theory in curved space-times (i.e., from “below”). A second problem involves matter couplings. Rovelli and Smolin (1993) have considered the coupling to a scalar field while Matschull and Nicoli (1993a, b) and Morales-Técolt and Rovelli (1994) have discussed fermion coupling. As I mentioned in section 4.3, Rovelli and Smolin use the scalar field to define an approximate time variable and thus recover a time dependent Schrödinger equation with a true Hamiltonian. For spinor field sources, they are developing an approximation technique to analyze the action of this Hamiltonian. A great deal of physical understanding is expected to come from such analyses of matter couplings.

Different people, of course, have different expectations of quantum gravity. To conclude, I would like to ask your indulgence for a moment or two as I express my personal views and prejudices.

The aim of the canonical program, as I see it, is to find out if a consistent quantum version of general relativity coupled to matter fields can exist non-perturbatively. Thus, one focusses on certain issues and, at least at the first go, ignores others. The focus is on quantum geometry, on diffeomorphism invariance, on internal clocks, on mathematical methods needed in any non-perturbative treatment of gravity, and, although these were not discussed at this workshop, on new approximation methods. The program is certainly far from being complete. However, it has not run into an unsurmountable obstacle either. The basic spirit, I believe, is the same as the one that drove physicists when they were groping in the dark during the development of quantum mechanics. One pushes the promising ideas and the plausible techniques as hard as one can. Either they keep working or break down at some point. If they break down, one focuses on where and why they fail. Occasionally, the breakdown suggests brand new strategies and changes the direction of the effort substantially. We know that, during the development of quantum mechanics, most ideas did not work and, at any given time, almost everyone was barking up the wrong tree. And yet this strategy of pushing the ideas ruthlessly as far as they can go always produced interesting results. In the same spirit, the hope now is that we will learn something interesting and perhaps even something deep. The evolution from metrics to connections to loops to weaves and knots represents progress along these lines.

Even if the program succeeds and produces a consistent, non-perturbative theory, there is no guarantee, of course, that this theory would be the correct one. Indeed, to make concrete progress, some issues were put aside at first and, in the final analysis, these may well turn out to be so central to the problem that they have to be incorporated right at the start. For example, it may well be, as the majority of the particle physics community holds, that real progress would not occur unless one has, from the very beginning, a principle which unifies the dynamics of all interactions. It may be inappropriate to seek insights into the quantum nature of geometry without incorporating *all* excitations of the superstring. It may also be, as several relativists believe, that real progress would not occur unless the approach is geared to tackle the measurement problems of quantum mechanics

from the start; unless the mathematical structure underlying quantum mechanics is made to absorb some fundamental non-linearity at the outset. Most people working on this canonical approach are sympathetic to these ideas in a general way. Indeed, work is in progress on both of these frontiers. However, the general sentiment is that the approach need not be *based* on such premises. Once the subject has evolved sufficiently, new ideas will come up to tackle such issues. There is often a tendency to underestimate the value of having a consistent, non-perturbative theory irrespective of whether it ultimately turns out to be the correct one physically. There appears to be an overriding sentiment that a quantum theory of gravity *must* solve all sorts of problems including those that are not, at least in any obvious way, intrinsic to gravity. And when one lists all these issues and all the associated problems, one is often so struck by the enormity of the task that a sense of hopelessness seems to take over. For a long time now, I think, there has been an undercurrent of pessimism at least in some parts of the relativity community: One does what one can but secretly (or, sometimes, openly!) believes that the task is *way* beyond us. This may well be the case. But I think it is also not obvious that this is really the case. Perhaps we should not try to solve all problems at once. We can suitably restrict our goals and pursue these “modest” programs with full enthusiasm and hope of success. I think we need a more outgoing attitude here, more aggressive spirits and more optimistic hearts!

### Acknowledgments

I benefited a great deal from the talks as well as informal discussions with the participants of this workshop. My special thanks go to Helmut Friedrich for organizing this stimulating meeting and for his patience with all administrative matters. This work was supported in part by the NSF grant PHY93-96246 and by the Eberly research fund of the Pennsylvania State University.

### References

- Achúcarro, A. & Townsend, P. (1986) Phys. Lett. **B180**, 85-89.
- Agishtein, A. & Migdal, A. (1992) Mod. Phys. Lett. **7**, 1039-1061.
- Amati, D. Ciafaloni, M. & Veneziano, G. (1990) Nucl. Phys. **B347**, 550-590.
- Aspinwall, P. S. (1993) Mirror symmetry, talk at the 1993 Rutherford-Appleton Meeting.
- Ashtekar, A. (1991) *Non-perturbative canonical gravity*, World Scientific, Singapore.
- Ashtekar, A. (1992) Mathematical problems of non-perturbative quantum general relativity, SU-GP-92/11-2; to appear in the proceedings of the 1992 Les Houches summer school, B. Julia (ed), Elsevier.
- Ashtekar, A., Husain, V., Rovelli, C., Samuel, J. & Smolin, L. (1989) Class. Quan. Grav. **6**, L185-L193.
- Ashtekar, A., Rovelli, C. & Smolin, L. (1991) Phys. Rev. **D44**, 1740-1755.
- Ashtekar, A. & Isham, C. (1992) Class. Quan. Grav. **9**, 1069-1100.
- Ashtekar, A. & Rovelli, C. (1992) Class. Quan. Grav. **91** 1121-1150.
- Ashtekar, A., Rovelli, C. & Smolin L. (1992) Phys. Rev. Lett. **69** 237-240.

- Ashtekar, A. & Varadarajan, M. (1992a,b) in preparation.
- Ashtekar, A. & Lewandowski, J. (1993a) Representation theory of analytic holonomy  $C^*$  algebras; to appear in *Knots and quantum gravity*, J. Baez (ed), Oxford University press.
- Ashtekar A. & Lewandowski, J. (1993b) Differential calculus on  $\mathcal{A}/\mathcal{G}$ , in preparation.
- Ashtekar, A. & Loll, R. (1994) New loop representations for 2+1-dimensional gravity.
- Baez, J. (1993a) Diffeomorphism-invariant generalized measures on the space of connections modulo gauge transformations, hep-th/9305045; to appear in the Proceedings of conference on quantum topology, L. Crane and D. Yetter (eds).
- Baez J. (1993b) Link invariants, functional integration and holonomy algebras, hep-th/9301063.
- Barbero-Gonzalez, J. F. (1993) General relativity as a theory of two connections, CGPG-93/9-5; Int. J. Mod. Phys. **D**(in press).
- Beig, R. (1994) article in this volume.
- Brügmann, B. (1994) article in this volume.
- Brügmann, B., Gambini, R. & Pullin, J. (1992a) Phys. Rev. Lett. **68**, 431-434.
- Brügmann, B., Gambini, R. & Pullin, J. (1992b) In: *XXth International conference on differential geometric methods in physics*, S. Cotta & A. Rocha (eds), World Scientific, Singapore.
- Brügmann, B. & Marinari, E. (1993) Phys. Rev. Lett. **70** 1908-1911.
- Brügmann, B. & Pullin, J. (1993) Nucl. Phys. **390** 399-438.
- Carlip, S. (1990) Phys. Rev. **D42**, 2647-2654.
- Carlip, S. (1993) Six ways to quantize (2+1)-dimensional gravity, gr-qc/9305020.
- de Wit, B., Matschull, H.J. & Nicolai, H. (1993) Phys. Lett. **B318**, 115.
- Di Bortolo, R. Gambini & Griego, J. (1992) Comm. Math. Phys. (to appear)
- Gambini, R. & Trias A. (1986) Nucl. Phys. **B278**, 436-448.
- Gross, D. & Mende, P (1988) Nucl. Phys. **B303**, 407-454.
- Guilliani, D. (1994) article in this volume.
- Hajiček, P. (1994) Article in this volume.
- Heisenberg, W. (1963) Interview by T.S. Kuhn of February 25th.
- Hosoya, A. & Nakao, K. (1990) Class. Quan. Grav. **7** 163-176.
- Iwasaki, J. & Rovelli, C. (1993) Int. J. Mod. Phys. **D1** 533-558.
- Kuchař (1970) J. Math. Phys. **11** 3322-3334.
- Loll, R. (1992) Nucl. Phys. **B368**, 121-142.
- Loll, R. (1994) article in this volume.
- Marolf, D. (1993) Loop representations of 2+1 gravity on a torus, SUGP-93/3/1.
- Marolf D. & Mourão J. (1993) Carrier space of the Ashtekar-Lewandowski measure, in preparation.
- Mena-Marugán, G. A. (1993) Reality conditions for Lorentzian and Euclidean gravity in the Ashtekar formulation, CGPG-93/9-3; Int. J. Mod. Phys. **D** (in press).
- Matschull, H.J. & Nicolai, H. (1993) DESY 93-073; Nucl Phys. **B** (in press)
- Mitter, P.K. & Viallet C. (1981) Commun. Math. Phys. **79** 43-58.
- Moncrief, V. (1989) J. Math. Phys. **30** 2907-2914.
- Morales-Técotl, H.A. & Rovelli, C. (1994) University of Pittsburgh pre-print.



- Nelson, J. & Regge, T. (1989) **A4**, 2021-2030.
- Nicolai, H. & Matschull, H.J. (1993) *J. Geo. & Phys.* **11** 15-62.
- Pais, A. (1987) In: *Paul Adrian Maurice Dirac*, B. N. Kursunoglu & E. P. Wigner (eds), Cambridge University Press, Cambridge, page 96.
- Pullin, J. (1993) Knot theory and quantum gravity in loop space: A premier, hep-th/9301028, to appear in the proceedings of the Vth Mexican school on particles and fields, J.L. Lucio (ed), Word Scientific, Singapore.
- Rovelli, C. & Smolin, L. (1990) *Nucl. Phys.* **B331**, 80-152.
- Rovelli, C. & Smolin, L. (1993) The physical Hamiltonian in non-perturbative quantum gravity, CGPC-93/8-2.
- Smolin, L. (1993) In: *General relativity and gravitation 1992*, R.J. Gleiser et al (eds), Institute of Physics, Bristol, pages 229-261.
- Streater, R. & Wightman, A. (1964) *PCT, spin and statistics, and all that*, Benjamin, New York.
- Thiemann, T. (1994) article in this volume.
- Witten, E. (1988) *Nucl. Phys.* **B311**, 46-78.
- Zegwaard, J. (1993) *Phys. Lett.* **B378** 217-220.
- Zegwaard, J. (1994) *The loop representation of canonical quantum gravity and its interpretation* Ph.D. thesis, University of Utrecht.